arXiv:1101.4113v1 [math.RT] 21 Jan 2011

AUSLANDER-REITEN TRANSLATIONS IN MONOMORPHISM CATEGORIES

BAO-LIN XIONG PU ZHANG* YUE-HUI ZHANG

Abstract. We generalize Ringel and Schmidmeier's theory on the Auslander-Reiten translation of the submodule category $\mathcal{S}_2(A)$ to the monomorphism category $\mathcal{S}_n(A)$. As in the case of $n = 2$, $\mathcal{S}_n(A)$ has Auslander-Reiten sequences, and the Auslander-Reiten translation $\tau_{\mathcal{S}}$ of $\mathcal{S}_n(A)$ can be explicitly formulated via τ of A-mod. Furthermore, if A is a selfinjective algebra, we study the periodicity of $\tau_{\mathcal{S}}$ on the objects of $\mathcal{S}_n(A)$, and of the Serre functor $F_{\mathcal{S}}$ on the objects of the stable monomorphism category $\underline{\mathcal{S}_n(A)}$. In particular, $\tau_{\mathcal{S}}^{2m(n+1)}X \cong X$ for $X \in \mathcal{S}_n(\Lambda(m,t))$; and $F_{\mathcal{S}}^{m(n+1)}X \cong X$ for $X \in \underline{\mathcal{S}_n(\Lambda(m,t))}$, where $\Lambda(m,t)$, $m \geq 1$, $t \geq 2$, are the selfinjective Nakayama algebras.

Key words and phrases. monomorphism category, Auslander-Reiten translation, triangulated category, Serre functor

Introduction

Throughout this paper, $n \geq 2$ is an integer, A an Artin algebra, and A-mod the category of finitely generated left A-modules. Let $\mathcal{S}_n(A)$ denote the monomorphism category of A (it is usually called the submodule category if $n = 2$).

The study of such a category goes back to G. Birkhoff [B], in which he initiates to classify the indecomposable objects of $\mathcal{S}_2(\mathbb{Z}/\langle p^t \rangle)$ (see also [RW]). In [Ar], $\mathcal{S}_n(R)$ is denoted by $\mathcal{C}(n,R)$, where R is a commutative uniserial ring; the complete list of $\mathcal{C}(n,R)$ of finite type, and of the representation types of $\mathcal{C}(n,k[x]/\langle x^t \rangle)$, are given by D. Simson [S] (see also [SW]). Recently, after the deep and systematic work of C. M. Ringel and M. Schmidmeier ([RS1] - [RS3]), the monomorphism category receives more attention. X. W. Chen [C] shows that $\mathcal{S}_2(\mathcal{A})$ of a Frobenius abelian category $\mathcal{A}$ is a Frobenius exact category. D. Kussin, H. Lenzing, and H. Meltzer [KLM] establish a surprising link between the stable submodule category with the singularity theory via weighted projective lines of type $(2,3,p)$. In [Z], $\mathcal{S}_n(\mathcal{X})$ is studied for any full subcategory $\mathcal{X}$ of A-mod, and it is proved that for a cotilting A-module T, there is a cotilting $T_n(A)$-module $\mathbf{m}(T)$ such that $\mathcal{S}_n({}^{\perp}T) = {}^{\perp}\mathbf{m}(T)$, where $T_n(A) = \begin{pmatrix} A & A & \cdots & A \\ 0 & A & \cdots & A \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A \end{pmatrix}_{n\times n}$ is the upper triangular matrix algebra

---

*The corresponding author.

2010 Mathematical Subject Classification. 16G10, 16G70, 18E30.

Supported by the NSF of China (10725104), and STCSM (09XD1402500).

of A, and ${}^{\perp}T$ is the left perpendicular category of T. As a consequence, for a Gorenstein algebra A, $\mathcal{S}_n({}^{\perp}A)$ is exactly the category of Gorenstein-projective $T_n(A)$-modules.

Ringel and Schmidmeier construct minimal monomorphisms, and then prove that $\mathcal{S}_2(A)$ is functorially finite in $T_2(A)$-mod. As a result, $\mathcal{S}_2(A)$ has Auslander-Reiten sequences. Surprisingly, the Auslander-Reiten translation $\tau_{\mathcal{S}}$ of $\mathcal{S}_2(A)$ can be explicitly formulated as $\tau_{\mathcal{S}}X \cong \mathbf{Mimo}\ \tau\ \mathbf{Cok}X$ for $X \in \mathcal{S}_2(A)$ ([RS2], Theorem 5.1), where τ is the Auslander-Reiten translation of A-mod. Applying this to selfinjective algebras, among others they get $\tau_{\mathcal{S}}^6 X \cong X$ for indecomposable nonprojective object $X \in \mathcal{S}_2(A)$, where A is a commutative uniserial algebra.

A beautiful theory should have a general version. The aim of this paper is to generalize Ringel and Schmidmeier's work on $\mathcal{S}_2(A)$ to $\mathcal{S}_n(A)$. As in the case of $n = 2$, $\mathcal{S}_n(A)$ has Auslander-Reiten sequences, and $\tau_{\mathcal{S}}$ of $\mathcal{S}_n(A)$ can be formulated in the same form as above: these can be achieved by using the idea in [RS2]. For selfinjective algebras, Sections 3 and 4 of this paper contain new considerations. In order to express the higher power of $\tau_{\mathcal{S}}$, we need the concept of a rotation of an object in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$, which is defined in [RS2] for $n = 2$. In the general case, such a suitable definition needs to be chosen from different possibilities, and difficulties need to be overcome to justify that it is well-defined. Also, the Octahedral Axiom is needed in computing the higher power of the rotations, which is the key step in studying the periodicity of $\tau_{\mathcal{S}}$ and the Serre functor on the objects.

We outline this paper. In Section 1 we set up some basic properties of the categories $\mathrm{Mor}_n(A)$, $\mathcal{S}_n(A)$ and $\mathcal{F}_n(A)$, and of the functors $\mathbf{m}_i$, $\mathbf{p}_i$, $\mathbf{Ker}$, $\mathbf{Cok}$, $\mathbf{Mono}$, $\mathbf{Epi}$; and the construction of $\mathbf{Mimo}$. Section 2 is to transfer the Auslander-Reiten sequences of $\mathrm{Mor}_n(A)$ to those of $\mathcal{S}_n(A)$; and to give a formula for $\tau_{\mathcal{S}}$ of $\mathcal{S}_n(A)$ via τ of A-mod (Theorem 2.4).

In Section 3, A is a selfinjective algebra, and hence the stable category A-$\underline{\mathrm{mod}}$ is a triangulated category ([H]), and τ is a triangle functor of A-$\underline{\mathrm{mod}}$. Using the rotation and the Octahedral Axiom, we get a formula for $\underline{\tau_{\mathcal{S}}^{j(n+1)}X} \in \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ for $X \in \mathcal{S}_n(A)$ and $j \geq 1$ (Theorem 3.4). This can be applied to the study of the periodicity of $\tau_{\mathcal{S}}$ on objects. In particular, $\tau_{\mathcal{S}}^{2m(n+1)}X \cong X$ for $X \in \mathcal{S}_n(\Lambda(m,t))$ (Corollary 3.6), where $\Lambda(m,t),\ m \geq 1,\ t \geq 2$, are the selfinjective Nakayama algebras.

In Section 4, A is a finite-dimensional self-injective algebra over a field. By [Z], $\mathcal{S}_n(A)$ is exactly the category of Gorenstein projective $T_n(A)$-modules, and hence the stable monomorphism category $\underline{\mathcal{S}_n(A)}$ is a Hom-finite Krull-Schmidt triangulated category with Auslander-Reiten triangles. By Theorem I.2.4 of I. Reiten and M. Van den Bergh [RV], $\underline{\mathcal{S}_n(A)}$ has a Serre functor $F_{\mathcal{S}}$. We study the periodicity of $F_{\mathcal{S}}$ on the objects of $\underline{\mathcal{S}_n(A)}$ (Theorem 4.3). In particular, $F_{\mathcal{S}}^{m(n+1)}X \cong X$ for $X \in \underline{\mathcal{S}_n(\Lambda(m,t))}$ (Corollary 4.4).

In order to make the main clue clearer, we put the proofs of Lemmas 1.5 and 1.6 in Appendix 1.

Note that $\mathcal{S}_{n,2}$, $\mathcal{S}_{2,3}$, $\mathcal{S}_{2,4}$, $\mathcal{S}_{2,5}$, $\mathcal{S}_{3,3}$ and $\mathcal{S}_{4,3}$ are the only representation-finite cases among all $\mathcal{S}_{n,t} = \mathcal{S}_n(k[x]/\langle x^t\rangle),\ n \geq 2,\ t \geq 2$ ([S], Theorems 5.2 and 5.5). The Auslander-Reiten quivers of $\mathcal{S}_{2,t}$ with $t = 2, 3, 4, 5$ are given in [RS3]. In Appendix 2, we give the remaining cases. We also include the AR quivers of $\mathcal{S}_n(\Lambda(2,2))$ with $n = 3$ and 4.

1. **Basics of morphism categories**

We set up some basic properties of several categories and functors, which will be used throughout this paper.

1.1. An object of *the morphism category* $\mathrm{Mor}_n(A)$ is $X_{(\phi_i)} = \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix}_{(\phi_i)}$, where $\phi_i : X_{i+1} \to X_i$ are A-maps for $1 \leq i \leq n-1$; and a morphism $f = (f_i) : X_{(\phi_i)} \to Y_{(\psi_i)}$ is $\begin{pmatrix} f_1 \\ \vdots \\ f_n \end{pmatrix}$, where $f_i : X_i \to Y_i$ are A-maps for $1 \leq i \leq n$, such that the following diagram commutes

$$\begin{tikzcd}
X_n \arrow[r, "\phi_{n-1}"] \arrow[d, "f_n"'] & X_{n-1} \arrow[r] \arrow[d, "f_{n-1}"] & \cdots \arrow[r] & X_2 \arrow[r, "\phi_1"] \arrow[d, "f_2"] & X_1 \arrow[d, "f_1"] \\
Y_n \arrow[r, "\psi_{n-1}"] & Y_{n-1} \arrow[r] & \cdots \arrow[r] & Y_2 \arrow[r, "\psi_1"] & Y_1.
\end{tikzcd} \tag{1.1}$$

We call X_i the i-th branch of $X_{(\phi_i)}$, and ϕ_i the i-th morphism of $X_{(\phi_i)}$. It is well-known that $\mathrm{Mor}_n(A)$ is equivalent to $T_n(A)$-mod (see e.g. [Z], 1.4). Let $Z_{(\theta_i)} \xrightarrow{f} Y_{(\psi_i)} \xrightarrow{g} X_{(\phi_i)}$ be a sequence in $\mathrm{Mor}_n(A)$. Then it is exact at $Y_{(\psi_i)}$ if and only if each sequence $Z_i \xrightarrow{f_i} Y_i \xrightarrow{g_i} X_i$ in A-mod is exact at Y_i for each $1 \leq i \leq n$.

By definition, *the monomorphism category* $\mathcal{S}_n(A)$ is the full subcategory of $\mathrm{Mor}_n(A)$ consisting of the objects $X_{(\phi_i)}$, where $\phi_i : X_{i+1} \to X_i$ are monomorphisms for $1 \leq i \leq n-1$. Dually, *the epimorphism category* $\mathcal{F}_n(A)$ is the full subcategory of $\mathrm{Mor}_n(A)$ consisting of the objects $X_{(\phi_i)}$, where $\phi_i : X_{i+1} \to X_i$ are epimorphisms for $1 \leq i \leq n-1$. Since $\mathcal{S}_n(A)$ and $\mathcal{F}_n(A)$ are closed under direct summands and extensions, it follows that they are exact Krull-Schmidt categories, with the exact structure in $\mathrm{Mor}_n(A)$.

The kernel functor $\mathbf{Ker} : \mathrm{Mor}_n(A) \to \mathcal{S}_n(A)$ is given by

$$\begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_{n-1} \\ X_n \end{pmatrix}_{(\phi_i)} \mapsto \begin{pmatrix} X_n \\ \mathrm{Ker}(\phi_1 \cdots \phi_{n-1}) \\ \vdots \\ \mathrm{Ker}(\phi_{n-2}\phi_{n-1}) \\ \mathrm{Ker}\,\phi_{n-1} \end{pmatrix}_{(\phi'_i)},$$

where $\phi'_1 : \mathrm{Ker}(\phi_1 \cdots \phi_{n-1}) \hookrightarrow X_n$, and $\phi'_i : \mathrm{Ker}(\phi_i \cdots \phi_{n-1}) \hookrightarrow \mathrm{Ker}(\phi_{i-1} \cdots \phi_{n-1}),\ 2 \leq i \leq n-1$, are the canonical monomorphisms. For a morphism $f : X \to Y$ in $\mathrm{Mor}_n(A)$, $\mathbf{Ker}f : \mathbf{Ker}X \to \mathbf{Ker}Y$ is naturally defined via a commutative diagram induced from

(1.1). We also need *the cokernel functor* $\mathbf{Cok}: \operatorname{Mor}_n(A) \to \mathcal{F}_n(A)$ given by

$$\begin{pmatrix} X_1\\ X_2\\ \vdots\\ X_{n-1}\\ X_n\end{pmatrix}_{(\phi_i)} \mapsto \begin{pmatrix}\operatorname{Coker}\phi_1\\ \operatorname{Coker}(\phi_1\phi_2)\\ \vdots\\ \operatorname{Coker}(\phi_1\cdots\phi_{n-1})\\ X_1\end{pmatrix}_{(\phi_i'')},$$

where $\phi_i'': \operatorname{Coker}(\phi_1\cdots\phi_{i+1}) \twoheadrightarrow \operatorname{Coker}(\phi_1\cdots\phi_i)$, $1 \le i \le n-2$, and $\phi_{n-1}'': X_1 \twoheadrightarrow \operatorname{Coker}(\phi_1\cdots\phi_{n-1})$ are the canonical epimorphisms. It is clear that the restriction of the kernel functor $\mathbf{Ker}: \mathcal{F}_n(A) \to \mathcal{S}_n(A)$ is an equivalence with quasi-inverse the restriction of the cokernel functor $\mathbf{Cok}: \mathcal{S}_n(A) \to \mathcal{F}_n(A)$.

1.2. For each $1 \le i \le n$, the functors $\mathbf{m}_i: A\text{-mod}\to \mathcal{S}_n(A)$ and $\mathbf{p}_i: A\text{-mod}\to \mathcal{F}_n(A)$ are defined as follows. For $M \in A$-mod,

$$(\mathbf{m}_i(M))_j = \begin{cases} M, & 1 \le j \le i;\\ 0, & i+1 \le j \le n; \end{cases} \qquad (\mathbf{p}_i(M))_j = \begin{cases} 0, & 1 \le j \le n-i;\\ M, & n-i+1 \le j \le n. \end{cases}$$

The j-th morphism of $\mathbf{m}_i(M)$ is id_M if $1 \le j < i$, and 0 if $i \le j \le n-1$; and the j-th morphism of $\mathbf{p}_i(M)$ is 0 if $1 \le j < n-i+1$, and id_M if $n-i+1 \le j \le n-1$. For an A-map $f: M \to N$, define

$$\mathbf{m}_i(f) = \begin{pmatrix} f\\ \vdots\\ f\\ 0\\ \vdots\\ 0\end{pmatrix} : \begin{pmatrix} M\\ \vdots\\ M\\ 0\\ \vdots\\ 0\end{pmatrix} \to \begin{pmatrix} N\\ \vdots\\ N\\ 0\\ \vdots\\ 0\end{pmatrix}; \qquad \mathbf{p}_i(f) = \begin{pmatrix} 0\\ \vdots\\ 0\\ f\\ \vdots\\ f\end{pmatrix} : \begin{pmatrix} 0\\ \vdots\\ 0\\ M\\ \vdots\\ M\end{pmatrix} \to \begin{pmatrix} 0\\ \vdots\\ 0\\ N\\ \vdots\\ N\end{pmatrix}.$$

We have

$$\mathbf{Ker}\ \mathbf{p}_i(M) = \mathbf{m}_{n-i+1}(M), \qquad \mathbf{Cok}\ \mathbf{m}_i(M) = \mathbf{p}_{n-i+1}(M), \quad 1 \le i \le n. \tag{1.2}$$

Lemma 1.1. *(i) If P runs over all the indecomposable projective A-modules, then $\mathbf{m}_1(P), \cdots, \mathbf{m}_n(P)$ are the all indecomposable projective objects in $\operatorname{Mor}_n(A)$.*

(ii) If I runs over all the indecomposable injective A-modules, then $\mathbf{p}_1(I), \cdots, \mathbf{p}_n(I)$ are the all indecomposable injective objects in $\operatorname{Mor}_n(A)$.

(iii) The indecomposable projective objects in $\mathcal{S}_n(A)$ are exactly those in $\operatorname{Mor}_n(A)$.

(iv) If I runs over all the indecomposable injective A-modules, then $\mathbf{m}_1(I), \cdots, \mathbf{m}_n(I)$ are the all indecomposable injective objects in $\mathcal{S}_n(A)$.

(v) If P runs over all the indecomposable projective A-modules, then $\mathbf{p}_1(P), \cdots, \mathbf{p}_n(P)$ are the all indecomposable projective objects in $\mathcal{F}_n(A)$.

(vi) The indecomposable injective objects in $\mathcal{F}_n(A)$ are exactly those in $\operatorname{Mor}_n(A)$.

(vii) Let $\mathcal{N}_{\mathcal{M}}$ and $\mathcal{N}$ be the Nakayama functor of $\operatorname{Mor}_n(A)$ and of A-mod, respectively. Then for a projective A-module P, $\mathcal{N}_{\mathcal{M}}\mathbf{m}_i(P) = \mathbf{p}_{n-i+1}(\mathcal{N}P)$, $1 \le i \le n$.

Proof. For convenience, we include a justification. (i) can be seen from the equivalence $\mathrm{Mor}_n(A) \cong T_n(A)$-mod. For (ii), see e.g. Lemma 1.3(ii) in [Z]. (iii) follows from (i), and (vi) follows from (ii). Using the equivalence $\mathbf{Ker} : \mathcal{F}_n(A) \to \mathcal{S}_n(A)$ together with (vi) and (1.2), we see (iv). Using the equivalence $\mathbf{Cok} : \mathcal{S}_n(A) \to \mathcal{F}_n(A)$ together with (iii) and (1.2), we see (v). To see (vii), note that if P is indecomposable, then $\mathcal{N}_{\mathcal{M}}\mathbf{m}_i(P)$ is an indecomposable injective $T_n(A)$-module, hence by (ii) it is of the form $\mathbf{p}_j(I)$. Thus

$$\begin{pmatrix} 0 \\ \vdots \\ 0 \\ \mathrm{soc}(I) \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \mathrm{soc}(\mathbf{p}_j(I)) = \mathrm{soc}(\mathcal{N}_{\mathcal{M}}\mathbf{m}_i(P)) = \mathrm{top}(\mathbf{m}_i(P)) = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ \mathrm{top}(P) \\ 0 \\ \vdots \\ 0 \end{pmatrix}.$$

Thus $n - j + 1 = i, \ \mathrm{soc}(I) = \mathrm{top}(P)$, which means $\mathcal{N}_{\mathcal{M}}\mathbf{m}_i(P) = \mathbf{p}_{n-i+1}(\mathcal{N}P)$. ■

1.3. Recall the functors $\mathbf{Mono} : \mathrm{Mor}_n(A) \to \mathcal{S}_n(A)$ and $\mathbf{Epi} : \mathrm{Mor}_n(A) \to \mathcal{F}_n(A)$. The first one is given by

$$\begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_{n-1} \\ X_n \end{pmatrix}_{(\phi_i)} \mapsto \begin{pmatrix} X_1 \\ \operatorname{Im}\phi_1 \\ \vdots \\ \operatorname{Im}(\phi_1\cdots\phi_{n-2}) \\ \operatorname{Im}(\phi_1\cdots\phi_{n-1}) \end{pmatrix}_{(\phi'_i)},$$

where $\phi'_1 : \operatorname{Im}\phi_1 \hookrightarrow X_1$, and $\phi'_i : \operatorname{Im}(\phi_1 \cdots \phi_i) \hookrightarrow \operatorname{Im}(\phi_1 \cdots \phi_{i-1}), \ 2 \le i \le n-1$, are the canonical monomorphisms. The second one is given by

$$\begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_{n-1} \\ X_n \end{pmatrix}_{(\phi_i)} \mapsto \begin{pmatrix} \operatorname{Im}(\phi_1\cdots\phi_{n-1}) \\ \operatorname{Im}(\phi_2\cdots\phi_{n-1}) \\ \vdots \\ \operatorname{Im}\phi_{n-1} \\ X_n \end{pmatrix}_{(\phi''_i)},$$

where $\phi''_i : \operatorname{Im}(\phi_{i+1} \cdots \phi_{n-1}) \twoheadrightarrow \operatorname{Im}(\phi_i \cdots \phi_{n-1}), \ 1 \le i \le n-2$, and $\phi''_{n-1} : X_n \twoheadrightarrow \operatorname{Im}\phi_{n-1}$ are the canonical epimorphisms. Then

$$\mathbf{Epi} \cong \mathbf{Cok}\ \mathbf{Ker}, \quad \mathbf{Mono} \cong \mathbf{Ker}\ \mathbf{Cok}, \tag{1.3}$$

and hence $\mathbf{Mono}X \cong X$ for $X \in \mathcal{S}_n(A)$, and $\mathbf{Epi}Y \cong Y$ for $Y \in \mathcal{F}_n(A)$.

For an A-map $f : X \to Y$, denote the canonical A-maps $X \twoheadrightarrow \operatorname{Im} f$ and $\operatorname{Im} f \hookrightarrow Y$ by $\widetilde{f}$ and incl, respectively. The following lemma can be similarly proved as in [RS2] for $n = 2$.

Lemma 1.2. *Let* $X = X_{(\phi_i)} \in \mathrm{Mor}_n(A)$. *Then*

(i) *The morphism* $\begin{pmatrix} 1_{X_1} \\ \widetilde{\phi_1} \\ \vdots \\ \widetilde{\phi_1\cdots\phi_{n-1}} \end{pmatrix} : X \twoheadrightarrow \mathbf{Mono}X$ *is a left minimal approximation of* X *in* $\mathcal{S}_n(A)$.

(ii) The morphism $\begin{pmatrix} \mathrm{incl} \\ \vdots \\ \mathrm{incl} \\ 1_{X_n} \end{pmatrix}$: $\mathbf{Epi}X \hookrightarrow X$ *is a right minimal approximation of* X *in* $\mathcal{F}_n(A)$.

1.4. For $X_{(\phi_i)} \in \mathrm{Mor}_n(A)$, we define $\mathbf{Mimo}X_{(\phi_i)} \in \mathcal{S}_n(A)$ and $\mathbf{Mepi}X_{(\phi_i)} \in \mathcal{F}_n(A)$ as follows (see [Z]). For each $1 \le i \le n-1$, fix an injective envelope $e_i' : \mathrm{Ker}\,\phi_i \hookrightarrow \mathrm{IKer}\,\phi_i$. Then we have an A-map $e_i : X_{i+1} \to \mathrm{IKer}\,\phi_i$, which is an extension of e_i'. Define $\mathbf{Mimo}X_{(\phi_i)}$ to be

$$\begin{pmatrix} X_1 \oplus \mathrm{IKer}\,\phi_1 \oplus \cdots \oplus \mathrm{IKer}\,\phi_{n-1} \\ X_2 \oplus \mathrm{IKer}\,\phi_2 \oplus \cdots \oplus \mathrm{IKer}\,\phi_{n-1} \\ \vdots \\ X_{n-1} \oplus \mathrm{IKer}\,\phi_{n-1} \\ X_n \end{pmatrix}_{(\theta_i)}, \quad \text{where} \quad \theta_i = \begin{pmatrix} \phi_i & 0 & 0 & \cdots & 0 \\ e_i & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}_{(n-i+1)\times(n-i)}.$$

By construction $\mathbf{Mimo}X_{(\phi_i)} \in \mathcal{S}_n(A)$. Since $e_1, \cdots, e_{n-1}$ are not unique, we need to verify that $\mathbf{Mimo}X_{(\phi_i)}$ is well-defined. This can be seen from Lemma 1.3(i) below.

The object $\mathbf{Mepi}X_{(\phi_i)}$ is dually defined. Namely, for each $1 \le i \le n-1$, fix a projective cover $\pi_i' : \mathrm{PCoker}\,\phi_i \twoheadrightarrow \mathrm{Coker}\,\phi_i$, then we have an A-map $\pi_i : \mathrm{PCoker}\,\phi_i \to X_i$, which is a lift of π_i', and define $\mathbf{Mepi}X_{(\phi_i)} \in \mathcal{F}_n(A)$ to be

$$\begin{pmatrix} X_1 \\ X_2 \oplus \mathrm{PCoker}\,\phi_1 \\ \vdots \\ X_{n-1} \oplus \mathrm{PCoker}\,\phi_{n-2} \oplus \cdots \oplus \mathrm{PCoker}\,\phi_1 \\ X_n \oplus \mathrm{PCoker}\,\phi_{n-1} \oplus \cdots \oplus \mathrm{PCoker}\,\phi_1 \end{pmatrix}_{(\sigma_i)}, \quad \text{where} \quad \sigma_i = \begin{pmatrix} \phi_i & \pi_i & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 1 \end{pmatrix}_{i\times(i+1)}.$$

Remark. (i) $\mathbf{Mimo}X = X$ *for* $X \in \mathcal{S}_n(A)$, *and* $\mathbf{Mepi}Y = Y$ *for* $Y \in \mathcal{F}_n(A)$.

(ii) If each X_i *has no nonzero injective direct summands, then* $\mathbf{Mimo}X_{(\phi_i)}$ *has no nonzero injective direct summands in* $\mathcal{S}_n(A)$. *If each* X_i *has no nonzero projective direct summands, then* $\mathbf{Mepi}X_{(\phi_i)}$ *has no nonzero projective direct summands in* $\mathcal{F}_n(A)$. *These can be seen from Lemma 1.1(iv) and (v), respectively.*

Lemma 1.3. *Let* $X \in \mathrm{Mor}_n(A)$. *Then*

(i) The morphism $\begin{pmatrix} (1,0,\cdots,0) \\ \vdots \\ (1,0) \\ 1 \end{pmatrix}$: $\mathbf{Mimo}X \twoheadrightarrow X$ *is a right minimal approximation of* X *in* $\mathcal{S}_n(A)$.

(ii) The morphism $\begin{pmatrix} 1 \\ \binom{1}{0} \\ \vdots \\ \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \end{pmatrix}$: $X \hookrightarrow \mathbf{Mepi}X$ *is a left minimal approximation of* X *in* $\mathcal{F}_n(A)$.

For a proof of Lemma 1.3 we refer to [RS2] for $n = 2$, and to [Z] in general case. By Lemmas 1.2 and 1.3, and by Auslander and Smalø [AS], we get the following consequence

Corollary 1.4. *The subcategories* $\mathcal{S}_n(A)$ *and* $\mathcal{F}_n(A)$ *are functorially finite in* $\mathrm{Mor}_n(A)$ *and hence have Auslander-Reiten sequences.*

This corollary is the starting point of this paper. From now on, denote by τ, $\tau_{\mathcal{M}}$, $\tau_{\mathcal{S}}$ and $\tau_{\mathcal{F}}$ the Auslander-Reiten translations of A-mod, $\mathrm{Mor}_n(A)$, $\mathcal{S}_n(A)$ and $\mathcal{F}_n(A)$, respectively.

1.5. Let $A\text{-}\underline{\mathrm{mod}}$ (resp. $A\text{-}\overline{\mathrm{mod}}$) denote the stable category of A-mod modulo projective A-modules (resp. injective A-modules). Then $\tau = \mathrm{DTr}$ induces an equivalence $A\text{-}\underline{\mathrm{mod}} \rightarrow A\text{-}\overline{\mathrm{mod}}$ with quasi-inverse $\tau^{-} = \mathrm{TrD}$ ([ARS], p.106). Let $\mathrm{Mor}_n(A\text{-}\overline{\mathrm{mod}})$ denote the morphism category of $A\text{-}\overline{\mathrm{mod}}$. Namely, an object of $\mathrm{Mor}_n(A\text{-}\overline{\mathrm{mod}})$ is $\overline{X_{(\phi_i)}} = X_{(\overline{\phi_i})} = \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix}_{(\overline{\phi_i})}$ with $\overline{\phi_i} : X_{i+1} \rightarrow X_i$ in $A\text{-}\overline{\mathrm{mod}}$ for $1 \leq i \leq n-1$; and a morphism from $X_{(\overline{\phi_i})}$ to $Y_{(\overline{\theta_i})}$ is $\begin{pmatrix} \overline{f_1} \\ \vdots \\ \overline{f_n} \end{pmatrix}$, such that the corresponding version of (1.1) commutes in $A\text{-}\overline{\mathrm{mod}}$. Similarly, one has the morphism category $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$, in which an object is denoted by $\underline{X_{(\phi_i)}} = X_{(\underline{\phi_i})}$.

The following two lemmas will be heavily used in Sections 2 and 3. In order to make the main clue clearer, we put their proofs in Appendix 1.

Lemma 1.5. *Let* $X_{(\phi_i)} \in \mathrm{Mor}_n(A)$.

(*i*) *Let* $I_2, \cdots, I_n$ *be injective* A*-modules such that* $X'_{(\phi'_i)} = \begin{pmatrix} X_1 \oplus I_2 \oplus \cdots \oplus I_n \\ \vdots \\ X_{n-1} \oplus I_n \\ X_n \end{pmatrix}_{(\phi'_i)} \in \mathcal{S}_n(A)$, *where each* ϕ'_i *is of the form* $\begin{pmatrix} \phi_i & * & \cdots & * \\ * & * & \cdots & * \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \cdots & * \end{pmatrix}_{(n-i+1)\times(n-i)}$. *Then* $X'_{(\phi'_i)} \cong \mathbf{Mimo}X_{(\phi_i)} \oplus J$, *where* J *is an injective object of* $\mathcal{S}_n(A)$.

(*ii*) *Let* $P_1, \cdots, P_{n-1}$ *be projective* A*-modules such that* $X''_{(\phi''_i)} = \begin{pmatrix} X_1 \\ X_2 \oplus P_1 \\ \vdots \\ X_n \oplus P_{n-1} \oplus \cdots \oplus P_1 \end{pmatrix}_{(\phi''_i)} \in \mathcal{F}_n(A)$, *where each* ϕ''_i *is of the form* $\begin{pmatrix} \phi_i & * & \cdots & * \\ * & * & \cdots & * \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \cdots & * \end{pmatrix}_{i\times(i+1)}$. *Then* $X''_{(\phi''_i)} \cong \mathbf{Mepi}X_{(\phi_i)} \oplus L$, *where* L *is a projective object of* $\mathcal{F}_n(A)$.

Lemma 1.6. *Let* $X_{(\phi_i)}$, $Y_{(\psi_i)} \in \mathrm{Mor}_n(A)$.

(*i*) *If all branches* X_i *and* Y_i *have no nonzero injective direct summands, then* $\mathbf{Mimo}X_{(\phi_i)} \cong \mathbf{Mimo}Y_{(\psi_i)}$ *in* $\mathcal{S}_n(A)$ *if and only if* $X_{(\overline{\phi_i})} \cong Y_{(\overline{\psi_i})}$ *in* $\mathrm{Mor}_n(A\text{-}\overline{\mathrm{mod}})$.

(ii) If all X_i *and* Y_i *have no nonzero projective direct summands, then* $\mathbf{Mepi}X_{(\phi_i)} \cong \mathbf{Mepi}Y_{(\psi_i)}$ *in* $\mathcal{F}_n(A)$ *if and only if* $X_{(\underline{\phi_i})} \cong Y_{(\underline{\psi_i})}$ *in* $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$*.*

2. The Auslander-Reiten translation of $\mathcal{S}_n(A)$

In this section, we first transfer the Auslander-Reiten sequences of $\mathrm{Mor}_n(A)$ to those of $\mathcal{S}_n(A)$ and $\mathcal{F}_n(A)$; and then give a formula of the Auslander-Reiten translation $\tau_{\mathcal{S}}$ of $\mathcal{S}_n(A)$ via τ of A-mod. Results and methods in this section are generalizations of the corresponding ones in the case of $n = 2$, due to Ringel and Schmidmeier [RS2].

2.1. The following fact is crucial for later use.

Lemma 2.1. *Let* $0 \to X_{(\phi_i)} \xrightarrow{f} Y \xrightarrow{g} Z \to 0$ *be an Auslander-Reiten sequence of* $\mathrm{Mor}_n(A)$*.*

(i) If $\mathbf{Ker}Z$ *is not projective, then* $0 \to \mathbf{Ker}X \xrightarrow{\mathbf{Ker}f} \mathbf{Ker}Y \xrightarrow{\mathbf{Ker}g} \mathbf{Ker}Z \to 0$ *is either split exact, or an Auslander-Reiten sequence of* $\mathcal{S}_n(A)$*.*

(ii) If $\mathbf{Cok}X$ *is not injective, then* $0 \to \mathbf{Cok}X \xrightarrow{\mathbf{Cok}f} \mathbf{Cok}Y \xrightarrow{\mathbf{Cok}g} \mathbf{Cok}Z \to 0$ *is either split exact, or an Auslander-Reiten sequence of* $\mathcal{F}_n(A)$*.*

Proof. We only prove (i). Put $g' = \mathbf{Ker}g$ and $f' = \mathbf{Ker}f$. By Snake Lemma, $0 \to \mathbf{Ker}X \xrightarrow{f'} \mathbf{Ker}Y \xrightarrow{g'} \mathbf{Ker}Z$ is exact. Assume that g' is not a split epimorphism. We claim that g' is right almost split. Let $v : W \to \mathbf{Ker}Z$ be a morphism in $\mathcal{S}_n(A)$ which is not a split epimorphism. Applying $\mathbf{Cok}$, we get $t' = \mathbf{Cok}v : \mathbf{Cok}W \to \mathbf{Cok}\ \mathbf{Ker}Z = \mathbf{Epi}Z$, which is not a split epimorphism, and hence the composition $t : \mathbf{Cok}W \xrightarrow{t'} \mathbf{Epi}Z \overset{\sigma}{\hookrightarrow} Z$ is not a split epimorphism. So, there is a morphism $s : \mathbf{Cok}W \to Y$ such that $t = gs$. Applying $\mathbf{Ker}$, we get $\mathbf{Ker}t = g'\mathbf{Ker}s$ with $\mathbf{Ker}s : W \to \mathbf{Ker}Y$. Since $\mathbf{Ker}\sigma = \mathrm{id}_{\mathbf{Ker}Z}$, we see $v = g'\mathbf{Ker}s$. This proves the claim.

Since g' is right almost split and $\mathbf{Ker}Z$ is not projective, it follows that g' is epic, and hence f' is not a split monomorphism. We claim that f' is left almost split. For this, let $p : \mathbf{Ker}X \to B$ be a morphism in $\mathcal{S}_n(A)$ which is not a split monomorphism. Take an injective envelope $(e_i) : \begin{pmatrix} B_1/B_2 \\ \vdots \\ B_1/B_n \end{pmatrix}_{(\pi_i)} \hookrightarrow \begin{pmatrix} I_1 \\ \vdots \\ I_{n-1} \end{pmatrix}_{(b'_i)}$ in $\mathrm{Mor}_{n-1}(A)$. Put $B' = \begin{pmatrix} I_1 \\ \vdots \\ I_{n-1} \\ B_1 \end{pmatrix}_{(b'_i)} \in \mathrm{Mor}_n(A)$, where b'_{n-1} is the composition $B_1 \overset{\pi_{n-1}}{\twoheadrightarrow} B_1/B_n \overset{e_{n-1}}{\hookrightarrow} I_{n-1}$. By construction we get a morphism $\begin{pmatrix} e_1 \\ \vdots \\ e_{n-1} \\ \mathrm{id}_{B_1} \end{pmatrix} : \mathbf{Cok}B \to B'$, and $\mathbf{Cok}B = \mathbf{Epi}B'$. Hence $\mathbf{Ker}B' = B$. Put $r' = (r'_i) = \mathbf{Cok}p : \mathbf{Epi}X \to \mathbf{Cok}B$. Then we have a morphism $\begin{pmatrix} e_1 r'_1 \\ \vdots \\ e_{n-1} r'_{n-1} \end{pmatrix}$:

$\begin{pmatrix} \operatorname{Im}(\phi_1\cdots\phi_{n-1}) \\ \operatorname{Im}(\phi_2\cdots\phi_{n-1}) \\ \vdots \\ \operatorname{Im}\phi_{n-1} \end{pmatrix} \to \begin{pmatrix} I_1 \\ \vdots \\ I_{n-1} \end{pmatrix}_{(b'_i)}$ in $\operatorname{Mor}_{n-1}(A)$. By the canonical monomorphism (j_i) : $\begin{pmatrix} \operatorname{Im}(\phi_1\cdots\phi_{n-1}) \\ \operatorname{Im}(\phi_2\cdots\phi_{n-1}) \\ \vdots \\ \operatorname{Im}\phi_{n-1} \end{pmatrix} \hookrightarrow \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_{n-1} \end{pmatrix}$ in $\operatorname{Mor}_{n-1}(A)$, we get a morphism $\begin{pmatrix} r_1 \\ \vdots \\ r_{n-1} \end{pmatrix} : \begin{pmatrix} X_1 \\ \vdots \\ X_{n-1} \end{pmatrix}_{(\phi_i)} \to \begin{pmatrix} I_1 \\ \vdots \\ I_{n-1} \end{pmatrix}_{(b'_i)}$ in $\operatorname{Mor}_{n-1}(A)$, such that $e_i r'_i = r_i j_i$ for $1 \leq i \leq n-1$. Then we get a morphism $r = (r_i) : X \to B'$ in $\operatorname{Mor}_n(A)$ by letting $r_n = r'_n$ (we only need to check $b'_{n-1} r_n = r_{n-1}\phi_{n-1}$: $b'_{n-1} r_n = b'_{n-1} r'_n = e_{n-1}\pi_{n-1} r'_n = e_{n-1} r'_{n-1}\phi'_{n-1} = r_{n-1} j_{n-1} \phi'_{n-1} = r_{n-1}\phi_{n-1}$). By construction we get $\mathbf{Epi}r = r' : \mathbf{Epi}X \to \mathbf{Epi}B' = \mathbf{Cok}B$. This process can be figured as follows.

$$\begin{tikzcd}
 & X_n \arrow[r, "r'_n"] \arrow[dl, "\mathrm{id}_{X_n}"'] \arrow[dd, two heads, "\phi'_{n-1}"] & B_1 \arrow[r, two heads, "\mathrm{id}_{B_1}"] \arrow[dd, two heads, "\pi_{n-1}"] & B_1 \arrow[dd, "b'_{n-1}"] \\
X_n \arrow[urrr, "r_n=r'_n"'] \arrow[dd, "\phi_{n-1}"'] & & & \\
 & \operatorname{Im}\phi_{n-1} \arrow[r, "r'_{n-1}"] \arrow[dl, hook', "j_{n-1}"'] \arrow[d, two heads, "\phi'_{n-2}"] & B_1/B_n \arrow[r, hook, "e_{n-1}"] \arrow[d, two heads, "\pi_{n-2}"] & I_{n-1} \arrow[d, "b'_{n-2}"] \\
X_{n-1} \arrow[urrr, dotted, "r_{n-1}"'] \arrow[d] & \vdots \arrow[d, two heads] & \vdots \arrow[d, two heads] & \vdots \arrow[d, "b'_2"] \\
\vdots \arrow[dd] & \operatorname{Im}(\phi_2\cdots\phi_{n-1}) \arrow[r, "r'_2"] \arrow[dl, hook', "j_2"'] \arrow[dd, two heads, "\phi'_1"] & B_1/B_3 \arrow[r, hook, "e_2"] \arrow[dd, two heads, "\pi_1"] & I_2 \arrow[dd, "b'_1"] \\
X_2 \arrow[urrr, dotted, "r_2"'] \arrow[dd, "\phi_1"'] & & & \\
 & \operatorname{Im}(\phi_1\cdots\phi_{n-1}) \arrow[r, "r'_1"] \arrow[dl, hook', "j_1"'] & B_1/B_2 \arrow[r, hook, "e_1"] & I_1 \\
X_1 \arrow[urrr, dotted, "r_1"'] & & &
\end{tikzcd}$$

Clearly, r is not a spit monomorphism (otherwise, $\mathbf{Epi}r = r' : \mathbf{Epi}X \to \mathbf{Epi}B' = \mathbf{Cok}B$ is a spit monomorphism, and hence $p = \mathbf{Ker}\ r'$ is a spit monomorphism). So there is a morphism $h : Y \to B'$ such that $r = hf : X \to B'$. Applying $\mathbf{Ker}$, we get $\mathbf{Ker}r = (\mathbf{Ker}h)f'$, where $\mathbf{Ker}h : \mathbf{Ker}Y \to \mathbf{Ker}B' = B$. Since $\mathbf{Ker}r = \mathbf{Ker}\ \mathbf{Epi}r = \mathbf{Ker}r' = \mathbf{Ker}\ \mathbf{Cok}p = p$, we get $p = (\mathbf{Ker}h)f'$. This proves the claim, and completes the proof. ■

Proposition 2.2. *Let* $0 \to X \overset{f}{\to} Y \overset{g}{\to} Z \to 0$ *be an Auslander-Reiten sequence of* $\operatorname{Mor}_n(A)$.

(*i*) *If* $Z \in \mathcal{F}_n(A)$, *and* Z *is not projective in* $\mathcal{F}_n(A)$, *then*

$$0 \to \mathbf{Epi}X \overset{\mathbf{Epi}f}{\to} \mathbf{Epi}Y \overset{\mathbf{Epi}g}{\to} Z \to 0 \tag{2.1}$$

is an Auslander-Reiten sequence of $\mathcal{F}_n(A)$*; and*

$$0 \to \mathbf{Ker}X \overset{\mathbf{Ker}f}{\to} \mathbf{Ker}Y \overset{\mathbf{Ker}g}{\to} \mathbf{Ker}Z \to 0 \tag{2.2}$$

is an Auslander-Reiten sequence of $\mathcal{S}_n(A)$.

(ii) *If* $X \in \mathcal{S}_n(A)$*, and* X *is not injective in* $\mathcal{S}_n(A)$*, then*

$$0 \to X \overset{\mathbf{Mono}f}{\longrightarrow} \mathbf{Mono}Y \overset{\mathbf{Mono}g}{\longrightarrow} \mathbf{Mono}Z \to 0$$

is an Auslander-Reiten sequence of $\mathcal{S}_n(A)$*; and*

$$0 \to \mathbf{Cok}X \overset{\mathbf{Cok}f}{\to} \mathbf{Cok}Y \overset{\mathbf{Cok}g}{\to} \mathbf{Cok}Z \to 0$$

is an Auslander-Reiten sequence of $\mathcal{F}_n(A)$.

Proof. We only show (i). Since $\mathbf{Ker} : \mathcal{F}_n(A) \to \mathcal{S}_n(A)$ is an equivalence, and Z is not projective in $\mathcal{F}_n(A)$, it follows that $\mathbf{Ker}Z$ is not projective, and hence by Lemma 2.1, (2.2) is either split exact, or an Auslander-Reiten sequence in $\mathcal{S}_n(A)$. Applying the equivalence $\mathbf{Cok} : \mathcal{S}_n(A) \to \mathcal{F}_n(A)$, and using $\mathbf{Epi} \cong \mathbf{Cok}\ \mathbf{Ker}$ and $Z \in \mathcal{F}_n(A)$, we see that (2.1) is either split exact, or an Auslander-Reiten sequence in $\mathcal{F}_n(A)$. While $\mathbf{Epi}Y \to Z$ is the composition of the canonical monomorphism $\mathbf{Epi}Y \hookrightarrow Y$ and the right almost split morphism $Y \to Z$, so $\mathbf{Epi}Y \to Z$ is not a split epimorphism, hence (2.1) is an Auslander-Reiten sequence in $\mathcal{F}_n(A)$, so is (2.2). ■

2.2. We have the following relationship between $\tau_{\mathcal{S}}$ and $\tau_{\mathcal{M}}$.

Corollary 2.3. (i) *If* $Z \in \mathcal{S}_n(A)$*, then* $\tau_{\mathcal{S}}Z \cong \mathbf{Ker}\ \tau_{\mathcal{M}}\ \mathbf{Cok}Z$*, and* $\tau_{\mathcal{S}}^{-}Z \cong \mathbf{Mono}\ \tau_{\mathcal{M}}^{-}Z$.

(ii) *If* $Z \in \mathcal{F}_n(A)$*, then* $\tau_{\mathcal{F}}Z \cong \mathbf{Epi}\ \tau_{\mathcal{M}}Z$*, and* $\tau_{\mathcal{F}}^{-}Z \cong \mathbf{Cok}\ \tau_{\mathcal{M}}^{-}\ \mathbf{Ker}Z$.

Proof. We only prove the first formula of (i). Assume that Z is indecomposable. If Z is projective, then $Z = \mathbf{m}_i(P)$ by Lemma 1.1(iii), where P is an indecomposable projective A-module. By the definition of $\tau_{\mathcal{M}}$ and a direct computation, we have $\tau_{\mathcal{M}}\mathbf{Cok}Z = \tau_{\mathcal{M}}\mathbf{Cok}\ \mathbf{m}_i(P) = \tau_{\mathcal{M}}\ \mathbf{p}_{n-i+1}(P) = \begin{pmatrix} * \\ \vdots \\ * \\ 0 \end{pmatrix}$, it follows that $\mathbf{Ker}\ \tau_{\mathcal{M}}\ \mathbf{Cok}Z = 0 = \tau_{\mathcal{S}}Z$. Assume that $Z \in \mathcal{S}_n(A)$ is not projective. Since $\mathbf{Cok} : \mathcal{S}_n(A) \to \mathcal{F}_n(A)$ is an equivalence, $\mathbf{Cok}Z \in \mathcal{F}_n(A)$ is not projective. By Lemma 1.1(i) and (v), $\mathbf{Cok}Z$ is an indecomposable nonprojective object in $\mathrm{Mor}_n(A)$. Replacing Z by $\mathbf{Cok}Z$ in (2.2), we get the assertion by $\mathbf{Ker}\ \mathbf{Cok}Z \cong Z$. ■

2.3. **Example.** Let k be a field, $A = k[x]/\langle x^2 \rangle$, and S be the simple A-module. Denote by $i : S \hookrightarrow A$ and $\pi : A \twoheadrightarrow S$ the canonical A-maps. Then we have the Auslander-Reiten sequence in $\mathrm{Mor}_3(A)$

$$0\longrightarrow\begin{pmatrix}0\\A\\S\end{pmatrix}_{(i,0)}\xrightarrow{\left(\begin{smallmatrix}\begin{pmatrix}0\\0\\1\end{pmatrix}\\ \begin{pmatrix}0\\ \binom{\pi}{1}\\ i\end{pmatrix}\end{smallmatrix}\right)}\begin{pmatrix}0\\0\\S\end{pmatrix}\oplus\begin{pmatrix}S\\S\oplus A\\A\end{pmatrix}_{(\binom{0}{1},(-1,\pi))}\xrightarrow{\left(\begin{pmatrix}0\\0\\-i\end{pmatrix},\begin{pmatrix}1\\(-1,\pi)\\1\end{pmatrix}\right)}\begin{pmatrix}S\\S\\A\end{pmatrix}_{(\pi,1)}\longrightarrow 0.$$

By (2.1), we get an Auslander-Reiten sequence in $\mathcal{F}_3(A)$

$$0\longrightarrow\begin{pmatrix}0\\S\\S\end{pmatrix}_{(1,0)}\xrightarrow{\left(\begin{smallmatrix}\begin{pmatrix}0\\0\\1\end{pmatrix}\\ \begin{pmatrix}0\\i\\i\end{pmatrix}\end{smallmatrix}\right)}\begin{pmatrix}0\\0\\S\end{pmatrix}\oplus\begin{pmatrix}S\\A\\A\end{pmatrix}_{(1,\pi)}\xrightarrow{\left(\begin{pmatrix}0\\0\\-i\end{pmatrix},\begin{pmatrix}1\\\pi\\1\end{pmatrix}\right)}\begin{pmatrix}S\\S\\A\end{pmatrix}_{(\pi,1)}\longrightarrow 0.$$

By (2.2), we get an Auslander-Reiten sequence in $\mathcal{S}_3(A)$

$$0\longrightarrow\begin{pmatrix}S\\S\\0\end{pmatrix}_{(0,1)}\xrightarrow{\left(\begin{smallmatrix}\begin{pmatrix}1\\1\\0\end{pmatrix}\\ \begin{pmatrix}i\\1\\0\end{pmatrix}\end{smallmatrix}\right)}\begin{pmatrix}S\\S\\S\end{pmatrix}_{(1,1)}\oplus\begin{pmatrix}A\\S\\0\end{pmatrix}_{(0,i)}\xrightarrow{\left(\begin{pmatrix}i\\1\\1\end{pmatrix},\begin{pmatrix}-1\\-1\\0\end{pmatrix}\right)}\begin{pmatrix}A\\S\\S\end{pmatrix}_{(1,i)}\longrightarrow 0.$$

2.4. In Corollary 2.3(i), $\tau_{\mathcal{S}}$ is formulated via $\tau_{\mathcal{M}}$ of $\mathrm{Mor}_n(A)$. However, $\tau_{\mathcal{M}}$ is usually more complicated than τ. The rest of this section is to give a formula of $\tau_{\mathcal{S}}$ via τ.

Before stating the main result, we need a notation. For $X_{(\underline{\phi_i})}\in\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$, define $\tau X_{(\underline{\phi_i})}=\begin{pmatrix}\tau X_1\\ \vdots\\ \tau X_n\end{pmatrix}_{(\tau\underline{\phi_i})}\in\mathrm{Mor}_n(A\text{-}\overline{\mathrm{mod}})$. Consider the full subcategory given by

$$\{Y_{(\psi_i)}=\begin{pmatrix}\tau X_1\\ \vdots\\ \tau X_n\end{pmatrix}_{(\psi_i)}\in\mathrm{Mor}_n(A)\mid Y_{\overline{(\psi_i)}}\cong\tau X_{(\underline{\phi_i})}\}.$$

Any object in this full subcategory will be denoted by $\tau X_{(\phi_i)}$ (we emphasize that this convention will cause no confusions). So, for $X_{(\phi_i)}\in\mathrm{Mor}_n(A)$ we have $\overline{\tau X_{(\phi_i)}}\cong\tau X_{(\underline{\phi_i})}$. By Lemma 1.6($i$), **Mimo** $\tau X_{(\phi_i)}$ is a well-defined object in $\mathcal{S}_n(A)$, and there are isomorphisms $\overline{\textbf{Mimo}\ \tau X_{(\phi_i)}}\cong\overline{\tau X_{(\phi_i)}}\cong\tau X_{(\underline{\phi_i})}$ in $\mathrm{Mor}_n(A\text{-}\overline{\mathrm{mod}})$. If A is selfinjective, then $\mathrm{Mor}_n(A\text{-}\overline{\mathrm{mod}})=\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$, so the isomorphism above is read as follows, which is needed in the next section

$$\underline{\textbf{Mimo}\ \tau X_{(\phi_i)}}\cong\underline{\tau X_{(\phi_i)}}\cong\tau X_{(\underline{\phi_i})}. \tag{2.3}$$

Similarly, one has the convention $\tau^-\ X_{(\phi_i)}$, and **Mepi** $\tau^-\ X_{(\phi_i)}\in\mathcal{F}_n(A)$ is well-defined.

The following result is a generalization of Theorem 5.1 of Ringel and Schmidmeier [RS2].

Theorem 2.4. *Let* $X_{(\phi_i)} \in \mathcal{S}_n(A)$*. Then*

(*i*) $\tau_{\mathcal{S}} X_{(\phi_i)} \cong \textbf{Mimo}\ \tau\ \textbf{Cok}X_{(\phi_i)}$.

(*ii*) $\tau_{\mathcal{S}}^{-} X_{(\phi_i)} \cong \textbf{Ker Mepi}\ \tau^{-}\ X_{(\phi_i)}$.

Proof. We only prove (*i*). Recall $\textbf{Cok}X_{(\phi_i)} = \begin{pmatrix} \operatorname{Coker}\phi_1 \\ \operatorname{Coker}(\phi_1\phi_2) \\ \vdots \\ \operatorname{Coker}(\phi_1\cdots\phi_{n-1}) \\ X_1 \end{pmatrix}_{(\phi'_i)}$. Fix a minimal projective presentation $Q_n \overset{d_n}{\to} P \overset{e}{\to} X_1 \to 0$. Then we get the following commutative diagram with exact rows

$$\begin{tikzcd}
Q_n \arrow[r, "d_n"] \arrow[d, dashed, "s_{n-1}"'] & P \arrow[r, "e"] \arrow[d, equal] & X_1 \arrow[r] \arrow[d, two heads, "\phi'_{n-1}"] & 0 \\
Q_{n-1} \arrow[r, "d_{n-1}"] \arrow[d, dashed] & P \arrow[r, "\phi'_{n-1}e"] \arrow[d, equal] & \operatorname{Coker}(\phi_1\cdots\phi_{n-1}) \arrow[r] \arrow[d, two heads] & 0 \\
\vdots \arrow[d, dashed, "s_1"'] & \vdots \arrow[d, equal] & \vdots \arrow[d, two heads, "\phi'_1"] & \\
Q_1 \arrow[r, "d_1"] & P \arrow[r, "\phi'_1\cdots\phi'_{n-1}e"] & \operatorname{Coker}(\phi_1) \arrow[r] & 0
\end{tikzcd} \tag{2.4}$$

where $Q_i \twoheadrightarrow \operatorname{Ker}(\phi'_i\cdots\phi'_{n-1}e)$ is a projective cover, and d_i is the composition $Q_i \twoheadrightarrow \operatorname{Ker}(\phi'_i\cdots\phi'_{n-1}e) \hookrightarrow P$. Applying the Nakayama functor $\mathcal{N} = D\operatorname{Hom}_A(-,\ {}_AA)$, we get the following commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \tau X_1 \arrow[r, "\sigma_n"] \arrow[d, dashed, "\alpha_{n-1}"'] & \mathcal{N}Q_n \arrow[r, "\mathcal{N}d_n"] \arrow[d, "\mathcal{N}s_{n-1}"'] & \mathcal{N}P \arrow[d, equal] \\
0 \arrow[r] & \tau\operatorname{Coker}(\phi_1\cdots\phi_{n-1}) \arrow[r, "\sigma_{n-1}"] \arrow[d, dashed, "\alpha_{n-2}"'] & \mathcal{N}Q_{n-1} \arrow[r, "\mathcal{N}d_{n-1}"] \arrow[d, "\mathcal{N}s_{n-2}"'] & \mathcal{N}P \arrow[d, equal] \\
 & \vdots \arrow[d, dashed] & \vdots \arrow[d] & \vdots \arrow[d, equal] \\
0 \arrow[r] & \tau\ \operatorname{Coker}(\phi_1\phi_2) \arrow[r, "\sigma_2"] \arrow[d, dashed, "\alpha_1"'] & \mathcal{N}Q_2 \arrow[r, "\mathcal{N}d_2"] \arrow[d, "\mathcal{N}s_1"'] & \mathcal{N}P \arrow[d, equal] \\
0 \arrow[r] & \tau\ \operatorname{Coker}\phi_1 \arrow[r, "\sigma_1"] & \mathcal{N}Q_1 \arrow[r, "\mathcal{N}d_1"] & \mathcal{N}P.
\end{tikzcd} \tag{2.5}$$

Step 1. By (2.4), we get a projective presentation

$$\bigoplus_{i=1}^{n} \mathbf{m}_i(Q_i) \xrightarrow{\left(\begin{pmatrix} d_1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \begin{pmatrix} d_2 \\ d_2 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \cdots, \begin{pmatrix} d_n \\ d_n \\ d_n \\ \vdots \\ d_n \end{pmatrix}\right)} \mathbf{m}_n(P) \xrightarrow{\begin{pmatrix} \phi'_1\cdots\phi'_{n-1}e \\ \vdots \\ \phi'_{n-1}e \\ e \end{pmatrix}} \textbf{Cok}X_{(\phi_i)} \longrightarrow 0 \tag{2.6}$$

(the exactness can be seen as follows: by (2.4) we have $\operatorname{Im} d_n \subseteq \operatorname{Im} d_{n-1} \subseteq \cdots \subseteq \operatorname{Im} d_1$, and hence $Q_i \oplus \cdots \oplus Q_n \overset{(d_i,\cdots,d_n)}{\longrightarrow} P \overset{\phi'_i\cdots\phi'_{n-1}e}{\longrightarrow} \operatorname{Coker}(\phi_i\cdots\phi_{n-1}) \to 0$ is exact). In order to obtain a minimal projective presentation from (2.6), we have to split off a direct summand of $\bigoplus\limits_{i=1}^{n}\mathbf{m}_i(Q_i)$. By Lemma 1.1($i$), this direct summand is of the form $\bigoplus\limits_{i=1}^{n-1}\mathbf{m}_i(Q'_i)$ where Q'_i is a direct summand of Q_i, $1\le i\le n-1$, since $\begin{pmatrix}\phi'_1\cdots\phi'_{n-1}e\\ \vdots\\ \phi'_{n-1}e\\ e\end{pmatrix}$ is already minimal and $Q_n\overset{d_n}{\to}P\overset{e}{\to}X_1\to 0$ is already a minimal projective presentation. Applying the Nakayama functor $\mathcal{N}_{\mathcal{M}}$, we get the exact sequence

$$0\longrightarrow\tau_{\mathcal{M}}\mathbf{Cok}X_{(\phi_i)}\oplus\mathcal{N}_{\mathcal{M}}(\bigoplus_{i=1}^{n-1}\mathbf{m}_i(Q'_i))\longrightarrow\mathcal{N}_{\mathcal{M}}(\bigoplus_{i=1}^{n}\mathbf{m}_i(Q_i))\longrightarrow\mathcal{N}_{\mathcal{M}}\mathbf{m}_n(P).$$

By Lemma 1.1(vii) this exact sequence can be written as

$$0\longrightarrow\tau_{\mathcal{M}}\mathbf{Cok}X_{(\phi_i)}\oplus\bigoplus_{i=1}^{n-1}\mathbf{p}_{n-i+1}(\mathcal{N}Q'_i)\longrightarrow\bigoplus_{i=1}^{n}\mathbf{p}_{n-i+1}(\mathcal{N}Q_i)\xrightarrow{d}\mathbf{p}_1(\mathcal{N}P)\tag{2.7}$$

where $d=\left(\begin{pmatrix}0\\ \vdots\\ 0\\ \mathcal{N}d_1\end{pmatrix},\cdots,\begin{pmatrix}0\\ \vdots\\ 0\\ \mathcal{N}d_n\end{pmatrix}\right)$.

Step 2. Write $Y_{(\theta_i)}=\tau_{\mathcal{M}}\mathbf{Cok}X_{(\phi_i)}\oplus\bigoplus\limits_{i=1}^{n-1}\mathbf{p}_{n-i+1}(\mathcal{N}Q'_i)$. By taking the i-th branches and the n-th branches of terms in (2.7), we get the following commutative diagram with exact rows

$$\begin{tikzcd}
0 \arrow[r] & Y_n \arrow[r, "\binom{a}{b}"] \arrow[d, "\theta_i\cdots\theta_{n-1}"'] & \bigoplus\limits_{j=1}^{i}\mathcal{N}Q_j\oplus\bigoplus\limits_{j=i+1}^{n}\mathcal{N}Q_j \arrow[rr, "((\mathcal{N}d_1{,}\cdots{,}\mathcal{N}d_i){,}(\mathcal{N}d_{i+1}{,}\cdots{,}\mathcal{N}d_n))"] \arrow[d, "(1{,}0)"] & & \mathcal{N}P \arrow[d] \\
0 \arrow[r] & Y_i \arrow[r, "\cong"] & \bigoplus\limits_{j=1}^{i}\mathcal{N}Q_j \arrow[rr] & & 0.
\end{tikzcd}$$

In particular, $Y_n=\operatorname{Ker}(\mathcal{N}d_1,\cdots,\mathcal{N}d_n)$. The upper exact sequence means that

$$\begin{tikzcd}
Y_n \arrow[r, "-b"] \arrow[d, "a"] & \bigoplus\limits_{j=i+1}^{n}\mathcal{N}Q_j \arrow[d, "(\mathcal{N}d_{i+1}{,}\cdots{,}\mathcal{N}d_n)"] \\
\bigoplus\limits_{j=1}^{i}\mathcal{N}Q_j \arrow[r, "(\mathcal{N}d_1{,}\cdots{,}\mathcal{N}d_i)"] & \mathcal{N}P
\end{tikzcd}$$

is a pull back square, for each $1\le i\le n-1$. It follows that

$$\operatorname{Ker}(\theta_i\cdots\theta_{n-1})=\operatorname{Ker}a=\operatorname{Ker}(\mathcal{N}d_{i+1},\cdots,\mathcal{N}d_n),\ 1\le i\le n-1,$$

and hence $\mathbf{Ker}Y_{(\theta_i)}=\begin{pmatrix}\operatorname{Ker}(\mathcal{N}d_1,\cdots,\mathcal{N}d_n)\\ \operatorname{Ker}(\mathcal{N}d_2,\cdots,\mathcal{N}d_n)\\ \vdots\\ \operatorname{Ker}(\mathcal{N}d_{n-1},\mathcal{N}d_n)\\ \operatorname{Ker}\mathcal{N}d_n\end{pmatrix}_{(\theta'_i)}$. We explicitly compute $\mathbf{Ker}Y_{(\theta_i)}$ below.

Step 3. By (2.5) we get the following commutative diagram with exact rows:

$$\begin{tikzcd}
0 \arrow[r] & \tau X_1 \arrow[r, "\gamma_n"] \arrow[d, "\beta_{n-1}"'] & \mathcal{N}Q_n \arrow[r, "\mathcal{N}d_n"] \arrow[d, "\left(\begin{smallmatrix}0\\1\end{smallmatrix}\right)"] & \mathcal{N}P \arrow[d, equal] \\
0 \arrow[r] & \tau\,\mathrm{Coker}(\phi_1\cdots\phi_{n-1})\oplus\mathcal{N}Q_n \arrow[r, "\gamma_{n-1}"] \arrow[d, "\beta_{n-2}"'] & \mathcal{N}Q_{n-1}\oplus\mathcal{N}Q_n \arrow[r, "{(\mathcal{N}d_{n-1},\ \mathcal{N}d_n)}"] \arrow[d, "\left(\begin{smallmatrix}0\\E_2\end{smallmatrix}\right)"] & \mathcal{N}P \arrow[d, equal] \\
 & \vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d, equal] \\
0 \arrow[r] & \tau\,\mathrm{Coker}(\phi_1\phi_2)\oplus\bigoplus\limits_{i=3}^{n}\mathcal{N}Q_i \arrow[r, "\gamma_2"] \arrow[d, "\beta_1"'] & \mathcal{N}Q_2\oplus\bigoplus\limits_{i=3}^{n}\mathcal{N}Q_i \arrow[r, "{(\mathcal{N}d_2,\cdots,\mathcal{N}d_n)}"] \arrow[d, "\left(\begin{smallmatrix}0\\E_{n-1}\end{smallmatrix}\right)"] & \mathcal{N}P \arrow[d, equal] \\
0 \arrow[r] & \tau\,\mathrm{Coker}\,\phi_1\oplus\bigoplus\limits_{i=2}^{n}\mathcal{N}Q_i \arrow[r, "\gamma_1"] & \mathcal{N}Q_1\oplus\bigoplus\limits_{i=2}^{n}\mathcal{N}Q_i \arrow[r, "{(\mathcal{N}d_1,\cdots,\mathcal{N}d_n)}"] & \mathcal{N}P
\end{tikzcd} \tag{2.8}$$

where E_i is the identity matrix, $\gamma_i = \begin{pmatrix} \sigma_i & (-\mathcal{N}s_i & -\mathcal{N}(s_is_{i+1}) & \cdots & -\mathcal{N}(s_i\cdots s_{n-1})) \\ 0 & & & E_{n-i} & \end{pmatrix}_{(n-i+1)\times(n-i+1)}$ for $1 \le i \le n$, $\beta_i = \begin{pmatrix} \alpha_i & (0 & 0 & \cdots & 0) \\ \sigma_{i+1} & (-\mathcal{N}s_{i+1} & -\mathcal{N}(s_{i+1}s_{i+2}) & \cdots & -\mathcal{N}(s_{i+1}\cdots s_{n-1})) \\ 0 & & & E_{n-i-1} & \end{pmatrix}_{(n-i+1)\times(n-i)}$ for $1 \le i \le n-1$. From (2.8) we see $\mathbf{Ker}Y_{(\theta_i)} \cong \begin{pmatrix} \tau\,\mathrm{Coker}\,\phi_1\oplus\bigoplus\limits_{j=2}^{n}\mathcal{N}Q_j \\ \tau\,\mathrm{Coker}(\phi_1\phi_2)\oplus\bigoplus\limits_{j=3}^{n}\mathcal{N}Q_j \\ \vdots \\ \tau\,\mathrm{Coker}(\phi_1\cdots\phi_{n-1})\oplus\mathcal{N}Q_n \\ \tau X_1 \end{pmatrix}_{(\beta_i)}$. Applying Lemma 1.5($i$), it is isomorphic to $\mathbf{Mimo}\ \tau\ \mathbf{Cok}X_{(\phi_i)} \oplus J$, where J is an injective object in $\mathcal{S}_n(A)$. Thus

$$\begin{aligned}\mathbf{Mimo}\quad \tau\ \mathbf{Cok}X_{(\phi_i)} \oplus J \cong \mathbf{Ker}Y_{(\theta_i)} &\overset{Def.}{\cong} \mathbf{Ker}\ \tau_{\mathcal{M}}\ \mathbf{Cok}X_{(\phi_i)} \oplus \mathbf{Ker}\Big(\bigoplus_{i=1}^{n-1}\mathbf{p}_{n-i+1}(\mathcal{N}Q_i')\Big) \\ &\overset{Cor.2.3}{\cong} \tau_{\mathcal{S}}X_{(\phi_i)} \oplus \mathbf{Ker}\Big(\bigoplus_{i=1}^{n-1}\mathbf{p}_{n-i+1}(\mathcal{N}Q_i')\Big) \overset{(1.2)}{\cong} \tau_{\mathcal{S}}X_{(\phi_i)} \oplus \bigoplus_{i=1}^{n-1}\mathbf{m}_i(\mathcal{N}Q_i').\end{aligned}$$

Since $\mathbf{Mimo}\ \tau\ \mathbf{Cok}X_{(\phi_i)}$ and $\tau_{\mathcal{S}}X_{(\phi_i)}$ have no nonzero injective direct summands in $\mathcal{S}_n(A)$ (cf. Remark (ii) in 1.4), and $\mathcal{S}_n(A)$ is Krull-Schmidt, we get $\tau_{\mathcal{S}}X_{(\phi_i)} \cong \mathbf{Mimo}\ \tau\ \mathbf{Cok}X_{(\phi_i)}$. ■

2.5. **Example.** Let A, S, i, and π be as in 2.3. Then there are 6 indecomposable non-projective objects in $\mathcal{S}_3(A)$. By Theorem 2.4 we have

$$\tau_{\mathcal{S}}\begin{pmatrix} A \\ S \\ 0 \end{pmatrix}_{(0,i)} = \mathbf{Mimo}\ \tau\begin{pmatrix} S \\ A \\ A \end{pmatrix}_{(1,\pi)} = \mathbf{Mimo}\begin{pmatrix} S \\ 0 \\ 0 \end{pmatrix}_{(0,0)} = \begin{pmatrix} S \\ 0 \\ 0 \end{pmatrix}_{(0,0)}$$

$$\tau_{\mathcal{S}}\begin{pmatrix} S \\ 0 \\ 0 \end{pmatrix}_{(0,0)} = \mathbf{Mimo}\ \tau\begin{pmatrix} S \\ S \\ S \end{pmatrix}_{(1,1)} = \mathbf{Mimo}\begin{pmatrix} S \\ S \\ S \end{pmatrix}_{(1,1)} = \begin{pmatrix} S \\ S \\ S \end{pmatrix}_{(1,1)}$$

$$\tau_{\mathcal{S}}\begin{pmatrix} S\\ S\\ S \end{pmatrix}_{(1,1)} = \mathbf{Mimo}\ \tau\begin{pmatrix} 0\\ 0\\ S \end{pmatrix}_{(0,0)} = \mathbf{Mimo}\begin{pmatrix} 0\\ 0\\ S \end{pmatrix}_{(0,0)} = \begin{pmatrix} A\\ A\\ S \end{pmatrix}_{(i,1)}$$

$$\tau_{\mathcal{S}}\begin{pmatrix} A\\ A\\ S \end{pmatrix}_{(i,1)} = \mathbf{Mimo}\ \tau\begin{pmatrix} 0\\ S\\ A \end{pmatrix}_{(\pi,0)} = \mathbf{Mimo}\begin{pmatrix} 0\\ S\\ 0 \end{pmatrix}_{(0,0)} = \begin{pmatrix} A\\ S\\ 0 \end{pmatrix}_{(1,i)}$$

$$\tau_{\mathcal{S}}\begin{pmatrix} S\\ S\\ 0 \end{pmatrix}_{(0,1)} = \mathbf{Mimo}\ \tau\begin{pmatrix} 0\\ S\\ S \end{pmatrix}_{(1,0)} = \mathbf{Mimo}\begin{pmatrix} 0\\ S\\ S \end{pmatrix}_{(1,0)} = \begin{pmatrix} A\\ S\\ S \end{pmatrix}_{(1,i)}$$

$$\tau_{\mathcal{S}}\begin{pmatrix} A\\ S\\ S \end{pmatrix}_{(1,i)} = \mathbf{Mimo}\ \tau\begin{pmatrix} S\\ S\\ A \end{pmatrix}_{(\pi,1)} = \mathbf{Mimo}\begin{pmatrix} S\\ S\\ 0 \end{pmatrix}_{(0,1)} = \begin{pmatrix} S\\ S\\ 0 \end{pmatrix}_{(0,1)}$$

The Auslander-Reiten quiver of $\mathcal{S}_3(A)$ looks like

$$\begin{tikzcd}
 & \begin{smallmatrix} A\\ 0\\ 0 \end{smallmatrix} \arrow[dr] & & \begin{smallmatrix} A\\ A\\ 0 \end{smallmatrix} \arrow[dr] & & \begin{smallmatrix} A\\ A\\ A \end{smallmatrix} \arrow[dr] & \\
\begin{smallmatrix} S\\ 0\\ 0 \end{smallmatrix} \arrow[ur] \arrow[dr] & & \begin{smallmatrix} A\\ S\\ 0 \end{smallmatrix} \arrow[ll, dotted] \arrow[ur] \arrow[dr] & & \begin{smallmatrix} A\\ A\\ S \end{smallmatrix} \arrow[ll, dotted] \arrow[ur] \arrow[dr] & & \begin{smallmatrix} S\\ S\\ S \end{smallmatrix} \arrow[ll, dotted] \\
 & \begin{smallmatrix} S\\ S\\ 0 \end{smallmatrix} \arrow[ur] \arrow[dr] & & \begin{smallmatrix} A\\ S\\ S \end{smallmatrix} \arrow[ll, dotted] \arrow[ur] \arrow[dr] & & \begin{smallmatrix} S\\ S\\ 0 \end{smallmatrix} \arrow[ll, dotted] \arrow[ur] & \\
 & & \begin{smallmatrix} S\\ S\\ S \end{smallmatrix} \arrow[ur] & & \begin{smallmatrix} S\\ 0\\ 0 \end{smallmatrix} \arrow[ll, dotted] \arrow[ur] & &
\end{tikzcd}$$

2.6. Denote by $\mathcal{S}_n(A)_{\mathcal{I}}$ the full subcategory of $\mathcal{S}_n(A)$ consisting of all the objects which have no nonzero injective direct summands of $\mathcal{S}_n(A)$. By Theorem 2.4, we have

Corollary 2.5. *Every object in $\mathcal{S}_n(A)_{\mathcal{I}}$ has the form* $\mathbf{Mimo}X$, *where each X_i has no nonzero injective direct summands.*

The following result will be used in the next section, whose proof is omitted, since it is the same as the case of $n = 2$ (see [RS2], Corollary 5.4).

Corollary 2.6. *The canonical functor $W : \mathcal{S}_n(A)_{\mathcal{I}} \to \mathrm{Mor}_n(A\text{-}\overline{\mathrm{mod}})$ given by $X_{(\phi_i)} \mapsto X_{(\overline{\phi_i})}$ is dense, preserves indecomposables, and reflects isomorphisms.*

2.7. Dual to Theorem 2.4 we have

Theorem 2.7. *Let $X \in \mathcal{F}_n(A)$. Then* $\tau_{\mathcal{F}}X \cong \mathbf{Cok}\ \mathbf{Mimo}\ \tau X$, *and* $\tau_{\mathcal{F}}^{-}X \cong \mathbf{Mepi}\ \tau^{-}\ \mathbf{Ker}X$.

3. Applications to selfinjective algebras

Throughout this section, A is a selfinjective algebra. Then A-$\overline{\mathrm{mod}} = A$-$\underline{\mathrm{mod}}$ is a triangulated category with the suspension functor Ω^{-1} ([H], p.16), where Ω^{-1} is the cosyzygy of A. The distinguished triangles of A-$\underline{\mathrm{mod}}$ are exactly triangles isomorphic to those given by all the short exact sequences in A-mod. Note that Ω and $\mathcal{N}$ commute and $\tau \cong \Omega^2\mathcal{N} \cong \mathcal{N}\Omega^2$ is an endo-equivalence of A-$\underline{\mathrm{mod}}$ ([ARS], p.126), and that τ is a triangle functor.

A rotation of an object in $\mathrm{Mor}_2(A\text{-}\underline{\mathrm{mod}})$ is introduced by Ringel and Schmidmeier [RS2]. The definition of a rotation of an object in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ needs new considerations. We have to take up pages to justify that it is well-defined. Then we get a formula for

$\underline{\tau_{\mathcal{S}}^{j}X} \in \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ for $X \in \mathcal{S}_n(A)$ and $j \geq 1$. This is applied to the study of the periodicity of $\tau_{\mathcal{S}}$ on the objects of $\mathcal{S}_n(A)$. In particular, for the selfinjective Nakayama algebras $\Lambda(m,t)$ we have $\tau_{\mathcal{S}}^{2m(n+1)}X \cong X$ for $X \in \mathcal{S}_n(\Lambda(m,t))$.

3.1. Let $X_{(\underline{\phi_i})} \in \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$. Just choose ϕ_i as representatives for the morphisms $\underline{\phi_i}$ in A-$\underline{\mathrm{mod}}$. Let $h_{i+1} : X_{i+1} \to I_{i+1}$ be an injective envelope with cokernel $\Omega^{-1}X_{i+1}$, $1 \leq i \leq n-1$. Taking pushout we get the following commutative diagram with exact rows

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "h_{i+1}"] \arrow[d, "\phi_1\cdots\phi_i"'] & I_{i+1} \arrow[r] \arrow[d, "j_{i+1}"] & \Omega^{-1}X_{i+1} \arrow[r] \arrow[d, equal] & 0 \\
0 \arrow[r] & X_1 \arrow[r, "g_{i+1}"] & Y_{i+1}^{1} \arrow[r] & \Omega^{-1}X_{i+1} \arrow[r] & 0.
\end{tikzcd}$$

This gives the exact sequence

$$0 \longrightarrow X_{i+1} \xrightarrow{\begin{pmatrix}\phi_1\cdots\phi_i \\ h_{i+1}\end{pmatrix}} X_1 \oplus I_{i+1} \xrightarrow{(g_{i+1},\ -j_{i+1})} Y_{i+1}^{1} \longrightarrow 0, \tag{3.1}$$

and induces the following commutative diagram with exact rows, $1 \leq i \leq n-2$

$$\begin{tikzcd}
0 \arrow[r] & X_{i+2} \arrow[r, "\begin{pmatrix}\phi_1\cdots\phi_{i+1} \\ h_{i+2}\end{pmatrix}"] \arrow[d, "\phi_{i+1}"'] & X_1 \oplus I_{i+2} \arrow[r, "{(g_{i+2},\ -j_{i+2})}"] \arrow[d, dashed, "\begin{pmatrix}1&0\\0&*\end{pmatrix}"] & Y_{i+2}^{1} \arrow[r] \arrow[d, dashed, "\psi_i"] & 0 \\
0 \arrow[r] & X_{i+1} \arrow[r, "\begin{pmatrix}\phi_1\cdots\phi_i \\ h_{i+1}\end{pmatrix}"] & X_1 \oplus I_{i+1} \arrow[r, "{(g_{i+1},\ -j_{i+1})}"] & Y_{i+1}^{1} \arrow[r] & 0.
\end{tikzcd} \tag{3.2}$$

By (3.2) we have $g_{i+1} = \psi_i g_{i+2}$, $1 \leq i \leq n-2$. Put $\psi_{n-1} = g_n$. Then $g_{i+1} = \psi_i \cdots \psi_{n-1}$, $1 \leq i \leq n-1$. By the construction of a distinguished triangle in A-$\underline{\mathrm{mod}}$, we get distinguished triangles from (3.1)

$$X_{i+1} \xrightarrow{\underline{\phi_1\cdots\phi_i}} X_1 \to Y_{i+1}^{1} \to \Omega^{-1}X_{i+1},\ 1 \leq i \leq n-1, \tag{3.3}$$

and by (3.2) we get the following commutative diagram, where the rows are distinguished triangles from (3.3)

$$\begin{tikzcd}
X_n \arrow[r] \arrow[d, "\underline{\phi_{n-1}}"] & X_1 \arrow[r, "\underline{\psi_{n-1}}"] \arrow[d, equal] & Y_n^{1} \arrow[r] \arrow[d, dashed, "\underline{\psi_{n-2}}"] & \Omega^{-1}X_n \arrow[d] \\
X_{n-1} \arrow[r] \arrow[d, "\underline{\phi_{n-2}}"] & X_1 \arrow[r] \arrow[d, equal] & Y_{n-1}^{1} \arrow[r] \arrow[d, dashed, "\underline{\psi_{n-3}}"] & \Omega^{-1}X_{n-1} \arrow[d] \\
\vdots \arrow[d, "\underline{\phi_2}"] & \vdots \arrow[d, equal] & \vdots \arrow[d, dashed, "\underline{\psi_1}"] & \vdots \arrow[d] \\
X_2 \arrow[r, "\underline{\phi_1}"] & X_1 \arrow[r] & Y_2^{1} \arrow[r] & \Omega^{-1}X_2.
\end{tikzcd} \tag{3.4}$$

The *rotation* $\mathbf{Rot}X_{(\underline{\phi_i})}$ of $X_{(\underline{\phi_i})}$ is defined to be

$$(X_1 \xrightarrow{\underline{\psi_{n-1}}} Y_n^{1} \longrightarrow \cdots \xrightarrow{\underline{\psi_1}} Y_2^{1}) \in \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$$

(here and in the following, for convenience we write the rotation in a row). We remark that $\mathbf{Rot}X_{\underline{(\phi_i)}}$ is well-defined: if $X_{\underline{(\phi_i)}} \cong Y_{\underline{(\theta_i)}}$ in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ with all X_i and Y_i having no nonzero injective direct summands, then $\mathbf{Mimo}X_{(\phi_i)} \cong \mathbf{Mimo}Y_{(\theta_i)}$ by Lemma 1.6(i), and hence $\mathbf{Rot}X_{\underline{(\phi_i)}} \cong \mathbf{Rot}Y_{\underline{(\theta_i)}}$, by Lemma 3.1 below.

Lemma 3.1. *Let* $X_{(\phi_i)} \in \mathrm{Mor}_n(A)$*. Then* $\mathbf{Rot}X_{\underline{(\phi_i)}} \cong \underline{\mathbf{Cok\ Mimo}X_{(\phi_i)}}$ *in* $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$.

Before proving Lemma 3.1, for later convenience, we restate Claim 2 in §4 of [RS2] in the more explicit way we will use.

Lemma 3.2. *Let* $0 \longrightarrow A \xrightarrow{\binom{f}{h}} B \oplus I \longrightarrow C \longrightarrow 0$ *be an exact sequence with* I *an injective* A*-module. Then there is an injective* A*-module* J *such that* $I = \mathrm{IKer}f \oplus J$*, and that the following diagram with exact rows commutes*

$$\begin{tikzcd}
0 \arrow[r] & A \arrow[r, "\left(\begin{smallmatrix} f \\ e \\ 0 \end{smallmatrix}\right)"] \arrow[d, equal] & B \oplus \mathrm{IKer}f \oplus J \arrow[r] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ w & 0 & 1 \end{smallmatrix}\right)"] & C' \oplus J \arrow[r] \arrow[d, "\wr"] & 0 \\
0 \arrow[r] & A \arrow[r, "\left(\begin{smallmatrix} f \\ e \\ h' \end{smallmatrix}\right)"] & B \oplus \mathrm{IKer}f \oplus J \arrow[r] & C \arrow[r] & 0,
\end{tikzcd}$$

where $h = \left(\begin{smallmatrix} e \\ h' \end{smallmatrix}\right)$*,* $e : A \to \mathrm{IKer}f$ *is an extension of the injective envelope* $\mathrm{Ker}\, f \hookrightarrow \mathrm{IKer}f$*,* $h' : A \to J$ *satisfies* $h'\,\mathrm{Ker}\, f = 0$*, and* $C' = \mathrm{Coker}\left(\begin{smallmatrix} f \\ e \end{smallmatrix}\right)$*.*

Proof of Lemma 3.1. We divide the proof into three steps.

Step 1. Recall $\mathbf{Mimo}X_{(\phi_i)} = \begin{pmatrix} X_1 \oplus \bigoplus\limits_{l=1}^{n-1} \mathrm{IKer}\phi_l \\ \vdots \\ X_{n-1} \oplus \mathrm{IKer}\,\phi_{n-1} \\ X_n \end{pmatrix}_{(\theta_i)}$ with $\theta_i = \begin{pmatrix} \phi_i & 0 & \cdots & 0 \\ e_i & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}_{(n-i+1)\times(n-i)}$,

$e_i : X_{i+1} \to \mathrm{IKer}\,\phi_i$ is an extension of the injective envelope $\mathrm{Ker}\,\phi_i \hookrightarrow \mathrm{IKer}\,\phi_i$, and

$$\mathbf{Cok\ Mimo}X_{(\phi_i)} = \begin{pmatrix} \mathrm{Coker}(\theta_1) \\ \mathrm{Coker}(\theta_1\theta_2) \\ \vdots \\ \mathrm{Coker}(\theta_1 \cdots \theta_{n-1}) \\ X_1 \oplus \bigoplus\limits_{l=1}^{n-1} \mathrm{IKer}\phi_l \end{pmatrix}_{(\theta'_i)}.$$

Since $\theta_1 \cdots \theta_i = diag(\alpha_i, E_{n-i-1}) : X_{i+1} \oplus \bigoplus\limits_{l=i+1}^{n-1} \mathrm{IKer}\phi_l \to X_1 \oplus \bigoplus\limits_{l=1}^{n-1} \mathrm{IKer}\phi_l$, where $\alpha_i = \begin{pmatrix} \phi_1 \cdots \phi_i \\ e_1\phi_2 \cdots \phi_i \\ \vdots \\ e_{i-1}\phi_i \\ e_i \end{pmatrix} : X_{i+1} \to X_1 \oplus \bigoplus\limits_{l=1}^{i} \mathrm{IKer}\phi_l$, and E_{n-i-1} is the identity matrix, we get the following commutative diagram with exact rows, $2 \le i \le n-1$,

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\alpha_i"] \arrow[d, "\phi_i"'] & X_1 \oplus \bigoplus\limits_{l=1}^{i} \mathrm{IKer}\phi_l \arrow[r, "\pi_i"] \arrow[d, "{(E_i,0)}"] & \mathrm{Coker}(\theta_1 \cdots \theta_i) \arrow[r] \arrow[d, "\theta'_{i-1}"] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\alpha_{i-1}"] & X_1 \oplus \bigoplus\limits_{l=1}^{i-1} \mathrm{IKer}\phi_l \arrow[r, "\pi_{i-1}"] & \mathrm{Coker}(\theta_1 \cdots \theta_{i-1}) \arrow[r] & 0
\end{tikzcd} \tag{3.5}$$

with $\pi_{n-1} = \theta'_{n-1}$.

Applying Lemma 3.2 to the upper exact sequence of (3.5) for $1 \le i \le n-1$, we get injective A-modules J_{i+1} such that $\bigoplus\limits_{l=1}^{i} \mathrm{IKer}\phi_l = \mathrm{IKer}(\phi_1 \cdots \phi_i) \oplus J_{i+1}$ and that the following diagram with exact rows commutes

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ 0 \end{smallmatrix}\right)"] \arrow[d, equal] & X_1 \oplus \mathrm{IKer}(\phi_1 \cdots \phi_i) \oplus J_{i+1} \arrow[r, "\left(\begin{smallmatrix} b_i^1 & b_i^2 & 0 \\ 0 & 0 & 1 \end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ w_i & 0 & 1 \end{smallmatrix}\right)"] & Z_{i+1} \oplus J_{i+1} \arrow[r] \arrow[d, "\beta_i", "\wr"'] & 0 \\
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ d_i \end{smallmatrix}\right)"] & X_1 \oplus \mathrm{IKer}(\phi_1 \cdots \phi_i) \oplus J_{i+1} \arrow[r, "\pi_i"] & \mathrm{Coker}(\theta_1 \cdots \theta_i) \arrow[r] & 0
\end{tikzcd} \tag{3.6}$$

where $\alpha_i = \left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ d_i \end{smallmatrix}\right)$, $a_i : X_{i+1} \to \mathrm{IKer}(\phi_1 \cdots \phi_i)$ is an extension of the injective envelope $\mathrm{Ker}(\phi_1 \cdots \phi_i) \hookrightarrow \mathrm{IKer}(\phi_1 \cdots \phi_i)$, $d_i : X_{i+1} \to J_{i+1}$ satisfies $d_i \,\mathrm{Ker}(\phi_1 \cdots \phi_i)$=0, and $Z_{i+1} = \mathrm{Coker}\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \end{smallmatrix}\right)$. Thus by (3.6) and (3.5) we get the following commutative diagram with exact rows for $2 \le i \le n-1$ (where the two rows in the middle come from (3.5)):

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ 0 \end{smallmatrix}\right)"] \arrow[d, equal] & X_1 \oplus \mathrm{IKer}(\phi_1 \cdots \phi_i) \oplus J_{i+1} \arrow[r, "\left(\begin{smallmatrix} b_i^1 & b_i^2 & 0 \\ 0 & 0 & 1 \end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ w_i & 0 & 1 \end{smallmatrix}\right)"] & Z_{i+1} \oplus J_{i+1} \arrow[r] \arrow[d, "\beta_i", "\wr"'] & 0 \\
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ d_i \end{smallmatrix}\right)"] \arrow[d, "\phi_i"'] & X_1 \oplus \mathrm{IKer}(\phi_1 \cdots \phi_i) \oplus J_{i+1} \arrow[r, "\pi_i"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & * & * \\ 0 & * & * \end{smallmatrix}\right)"] & \mathrm{Coker}(\theta_1 \cdots \theta_i) \arrow[r] \arrow[d, "\theta'_{i-1}"] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_{i-1} \\ a_{i-1} \\ d_{i-1} \end{smallmatrix}\right)"] \arrow[d, equal] & X_1 \oplus \mathrm{IKer}(\phi_1 \cdots \phi_{i-1}) \oplus J_i \arrow[r, "\pi_{i-1}"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ w_{i-1} & 0 & 1 \end{smallmatrix}\right)^{-1}"] & \mathrm{Coker}(\theta_1 \cdots \theta_{i-1}) \arrow[r] \arrow[d, "\beta_{i-1}^{-1}", "\wr"'] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_{i-1} \\ a_{i-1} \\ 0 \end{smallmatrix}\right)"] & X_1 \oplus \mathrm{IKer}(\phi_1 \cdots \phi_{i-1}) \oplus J_i \arrow[r, "\left(\begin{smallmatrix} b_{i-1}^1 & b_{i-1}^2 & 0 \\ 0 & 0 & 1 \end{smallmatrix}\right)"] & Z_i \oplus J_i \arrow[r] & 0.
\end{tikzcd}$$

Taking the first and the last rows, we get the following commutative diagram with exact rows:

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ 0 \end{smallmatrix}\right)"] \arrow[d, "\phi_i"'] & X_1 \oplus \mathrm{IKer}(\phi_1 \cdots \phi_i) \oplus J_{i+1} \arrow[r, "\left(\begin{smallmatrix} b_i^1 & b_i^2 & 0 \\ 0 & 0 & 1 \end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ c_{i1} & c_{i2} & * \\ * & * & * \end{smallmatrix}\right)"] & Z_{i+1} \oplus J_{i+1} \arrow[r] \arrow[d, "\beta_{i-1}^{-1}\theta'_{i-1}\beta_i"] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_{i-1} \\ a_{i-1} \\ 0 \end{smallmatrix}\right)"] & X_1 \oplus \mathrm{IKer}(\phi_1 \cdots \phi_{i-1}) \oplus J_i \arrow[r, "\left(\begin{smallmatrix} b_{i-1}^1 & b_{i-1}^2 & 0 \\ 0 & 0 & 1 \end{smallmatrix}\right)"] & Z_i \oplus J_i \arrow[r] & 0,
\end{tikzcd}$$

where for later convenience we write

$$\beta_{i-1}^{-1}\theta'_{i-1}\beta_i = \left(\begin{smallmatrix} f_{i-1} & * \\ * & * \end{smallmatrix}\right),\ 2 \le i \le n-1. \tag{3.7}$$

Taking off J_i and J_{i+1}, we get the following commutative diagram with exact rows, $2 \le i \le n-1$,

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \end{smallmatrix}\right)"] \arrow[d, "\phi_i"'] & X_1 \oplus \mathrm{IKer}(\phi_1\cdots\phi_i) \arrow[r, "(b_i^1,\ b_i^2)"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 \\ c_{i1} & c_{i2} \end{smallmatrix}\right)"] & Z_{i+1} \arrow[r] \arrow[d, "f_{i-1}"] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_{i-1} \\ a_{i-1} \end{smallmatrix}\right)"] & X_1 \oplus \mathrm{IKer}(\phi_1\cdots\phi_{i-1}) \arrow[r, "(b_{i-1}^1,\ b_{i-1}^2)"] & Z_i \arrow[r] & 0.
\end{tikzcd} \tag{3.8}$$

Step 2. Now we consider the rotation $\mathbf{Rot}X_{(\underline{\phi_i})}$. Recall from the beginning of this subsection that $h_{i+1} : X_{i+1} \to I_{i+1}$ is an injective envelope. For $1 \le i \le n-1$, applying Lemma 3.2 to (3.1), we get injective A-modules J'_{i+1} such that $I_{i+1} = \mathrm{IKer}(\phi_1\cdots\phi_i) \oplus J'_{i+1}$ and that the following diagram with exact rows commutes (cf. (3.6))

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ 0 \end{smallmatrix}\right)"] \arrow[d, equal] & X_1 \oplus \mathrm{IKer}(\phi_1\cdots\phi_i) \oplus J'_{i+1} \arrow[r, "\left(\begin{smallmatrix} b_i^1 & b_i^2 & 0 \\ 0 & 0 & 1 \end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ w'_i & 0 & 1 \end{smallmatrix}\right)"] & Z_{i+1} \oplus J'_{i+1} \arrow[r] \arrow[d, "\beta'_i", "\wr"'] & 0 \\
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ d'_i \end{smallmatrix}\right)"] & X_1 \oplus \mathrm{IKer}(\phi_1\cdots\phi_i) \oplus J'_{i+1} \arrow[r, "(g_{i+1},\ -j_{i+1})"] & Y_{i+1}^1 \arrow[r] & 0,
\end{tikzcd} \tag{3.9}$$

where $h_{i+1} = \left(\begin{smallmatrix} a_i \\ d'_i \end{smallmatrix}\right)$, and $d'_i : X_{i+1} \to J'_{i+1}$, satisfying $d'_i \,\mathrm{Ker}(\phi_1\cdots\phi_i) = 0$. Thus by (3.9) and (3.2) we get the following commutative diagram with exact rows for $2 \le i \le n-1$ (where the two rows in the middle come from (3.2)):

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ 0 \end{smallmatrix}\right)"] \arrow[d, equal] & X_1 \oplus \mathrm{IKer}(\phi_1\cdots\phi_i) \oplus J'_{i+1} \arrow[r, "\left(\begin{smallmatrix} b_i^1 & b_i^2 & 0 \\ 0 & 0 & 1 \end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ w'_i & 0 & 1 \end{smallmatrix}\right)"] & Z_{i+1} \oplus J'_{i+1} \arrow[r] \arrow[d, "\beta'_i", "\wr"'] & 0 \\
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_i \\ a_i \\ d'_i \end{smallmatrix}\right)"] \arrow[d, "\phi_i"'] & X_1 \oplus \mathrm{IKer}(\phi_1\cdots\phi_i) \oplus J'_{i+1} \arrow[r, "(g_{i+1},\ -j_{i+1})"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & * & * \\ 0 & * & * \end{smallmatrix}\right)"] & Y_{i+1}^1 \arrow[r] \arrow[d, "\psi_{i-1}"] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_{i-1} \\ a_{i-1} \\ d'_{i-1} \end{smallmatrix}\right)"] \arrow[d, equal] & X_1 \oplus \mathrm{IKer}(\phi_1\cdots\phi_{i-1}) \oplus J'_i \arrow[r, "(g_i,\ -j_i)"] \arrow[d, "\left(\begin{smallmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ w'_{i-1} & 0 & 1 \end{smallmatrix}\right)^{-1}"] & Y_i^1 \arrow[r] \arrow[d, "\beta'^{-1}_{i-1}", "\wr"'] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\left(\begin{smallmatrix} \phi_1\cdots\phi_{i-1} \\ a_{i-1} \\ 0 \end{smallmatrix}\right)"] & X_1 \oplus \mathrm{IKer}(\phi_1\cdots\phi_{i-1}) \oplus J'_i \arrow[r, "\left(\begin{smallmatrix} b_{i-1}^1 & b_{i-1}^2 & 0 \\ 0 & 0 & 1 \end{smallmatrix}\right)"] & Z_i \oplus J'_i \arrow[r] & 0.
\end{tikzcd}$$

Taking the first and the last rows, we get the following commutative diagram with exact rows:

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix}\phi_1\cdots\phi_i\\ a_i\\ 0\end{smallmatrix}\right)"] \arrow[d, "\phi_i"'] & X_1\oplus \mathrm{IKer}(\phi_1\cdots\phi_i)\oplus J'_{i+1} \arrow[r, "\left(\begin{smallmatrix}b_i^1 & b_i^2 & 0\\ 0 & 0 & 1\end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix}1 & 0 & 0\\ c'_{i1} & c'_{i2} & *\\ * & * & *\end{smallmatrix}\right)"] & Z_{i+1}\oplus J'_{i+1} \arrow[r] \arrow[d, "\beta'^{-1}_{i-1}\psi_{i-1}\beta'_i"] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\left(\begin{smallmatrix}\phi_1\cdots\phi_{i-1}\\ a_{i-1}\\ 0\end{smallmatrix}\right)"] & X_1\oplus \mathrm{IKer}(\phi_1\cdots\phi_{i-1})\oplus J'_i \arrow[r, "\left(\begin{smallmatrix}b_{i-1}^1 & b_{i-1}^2 & 0\\ 0 & 0 & 1\end{smallmatrix}\right)"] & Z_i\oplus J'_i \arrow[r] & 0,
\end{tikzcd}$$

where for later convenience we write

$$\beta'^{-1}_{i-1}\psi'_{i-1}\beta'_i = \left(\begin{smallmatrix}f'_{i-1} & *\\ * & *\end{smallmatrix}\right),\ 2\le i\le n-1. \tag{3.10}$$

Taking off J'_i and J'_{i+1}, we get the following commutative diagram with exact rows for $2\le i\le n-1$:

$$\begin{tikzcd}
0 \arrow[r] & X_{i+1} \arrow[r, "\left(\begin{smallmatrix}\phi_1\cdots\phi_i\\ a_i\end{smallmatrix}\right)"] \arrow[d, "\phi_i"'] & X_1\oplus \mathrm{IKer}(\phi_1\cdots\phi_i) \arrow[r, "{(b_i^1,\ b_i^2)}"] \arrow[d, "\left(\begin{smallmatrix}1 & 0\\ c'_{i1} & c'_{i2}\end{smallmatrix}\right)"] & Z_{i+1} \arrow[r] \arrow[d, "f'_{i-1}"] & 0 \\
0 \arrow[r] & X_i \arrow[r, "\left(\begin{smallmatrix}\phi_1\cdots\phi_{i-1}\\ a_{i-1}\end{smallmatrix}\right)"] & X_1\oplus \mathrm{IKer}(\phi_1\cdots\phi_{i-1}) \arrow[r, "{(b_{i-1}^1,\ b_{i-1}^2)}"] & Z_i \arrow[r] & 0.
\end{tikzcd}$$

Comparing the above diagram with (3.8), by a computation we easily see that $f_i - f'_i$ factors through an injective A-module for each $1\le i\le n-2$.

Step 3. Now we get the following diagram, where the first row can be considered as **Cok Mimo**$X_{(\phi_i)}$ (we identify $X_1\oplus\bigoplus\limits_{l=1}^{n-1}\mathrm{IKer}\phi_l$ with $X_1\oplus\mathrm{IKer}(\phi_1\cdots\phi_{n-1})\oplus J_n$; and identify $X_1\oplus I_n$ with $X_1\oplus\mathrm{IKer}(\phi_1\cdots\phi_{n-1})\oplus J'_n$):

$$\begin{tikzcd}
X_1\oplus\bigoplus\limits_{l=1}^{n-1}\mathrm{IKer}\phi_l \arrow[r, "\theta'_{n-1}"] \arrow[d, "\wr"', "\left(\begin{smallmatrix}1 & 0 & 0\\ 0 & 1 & 0\\ w_{n-1} & 0 & 1\end{smallmatrix}\right)^{-1}"] & \mathrm{Coker}(\theta_1\cdots\theta_{n-1}) \arrow[r, "\theta'_{n-2}"] \arrow[d, "\wr"', "\beta_{n-1}^{-1}"] & \cdots \arrow[r] & \mathrm{Coker}(\theta_1\theta_2) \arrow[r, "\theta'_1"] \arrow[d, "\wr"', "\beta_2^{-1}"] & \mathrm{Coker}\,\theta_1 \arrow[d, "\beta_1^{-1}"] \\
X_1\oplus\mathrm{IKer}(\phi_1\cdots\phi_{n-1})\oplus J_n \arrow[r, "\left(\begin{smallmatrix}b_{n-1}^1 & b_{n-1}^2 & 0\\ 0 & 0 & 1\end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix}1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 0\end{smallmatrix}\right)"] & Z_n\oplus J_n \arrow[r, "\left(\begin{smallmatrix}f_{n-2} & *\\ * & *\end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix}1 & 0\\ 0 & 0\end{smallmatrix}\right)"] & \cdots \arrow[r] & Z_3\oplus J_3 \arrow[r, "\left(\begin{smallmatrix}f_1 & *\\ * & *\end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix}1 & 0\\ 0 & 0\end{smallmatrix}\right)"'] & Z_2\oplus J_2 \arrow[d, "\left(\begin{smallmatrix}1 & 0\\ 0 & 0\end{smallmatrix}\right)"'] \\
X_1\oplus\mathrm{IKer}(\phi_1\cdots\phi_{n-1})\oplus J'_n \arrow[r, "\left(\begin{smallmatrix}b_{n-1}^1 & b_{n-1}^2 & 0\\ 0 & 0 & 1\end{smallmatrix}\right)"'] \arrow[d, "\wr"', "\left(\begin{smallmatrix}1 & 0 & 0\\ 0 & 1 & 0\\ w'_{n-1} & 0 & 1\end{smallmatrix}\right)"] & Z_n\oplus J'_n \arrow[r, "\left(\begin{smallmatrix}f'_{n-2} & *\\ * & *\end{smallmatrix}\right)"'] \arrow[d, "\wr"', "\beta'_{n-1}"] & \cdots \arrow[r] & Z_3\oplus J'_3 \arrow[r, "\left(\begin{smallmatrix}f'_1 & *\\ * & *\end{smallmatrix}\right)"'] \arrow[d, "\wr"', "\beta'_2"] & Z_2\oplus J'_2 \arrow[d, "\wr"', "\beta'_1"] \\
X_1\oplus I_n \arrow[r, "{(\psi_{n-1},-j_n)}"] & Y_n^1 \arrow[r, "\psi_{n-2}"] & \cdots \arrow[r] & Y_3^1 \arrow[r, "\psi_1"] & Y_2^1
\end{tikzcd}$$

(note that the squares in the first two rows commute in A-mod: the left square comes from (3.6); and the remaining commutative squares come from (3.7). Also, note that the squares in the last two rows commute in A-mod: the left square comes from (3.9); and the remaining commutative squares come from (3.10)). However, the squares in the middle may **not** commute in A-mod; and the point is that they commute in A-$\underline{\text{mod}}$, as we explain below.

Note that the left square in the middle commutes by a direct computation. Since $f_i - f_i'$ factors through an injective A-module, $1 \leq i \leq n-2$, we realize that the remaining $n-2$ squares in the middle commute in A-$\underline{\text{mod}}$. It follows that the above diagram commutes in A-$\underline{\text{mod}}$. It is clear that the vertical morphisms are isomorphisms in A-$\underline{\text{mod}}$. Regarding the above diagram in A-$\underline{\text{mod}}$, the first row is exactly $\underline{\mathbf{Cok\ Mimo}X_{(\phi_i)}}$, and the last row is exactly $\mathbf{Rot}X_{(\underline{\phi_i})}$. Thus, $\mathbf{Rot}X_{(\underline{\phi_i})} \cong \underline{\mathbf{Cok\ Mimo}X_{(\phi_i)}}$ in $\mathrm{Mor}_n(A\text{-}\underline{\text{mod}})$. This completes the proof. ■

3.2. Let $X_{(\phi_i)} \in \mathrm{Mor}_n(A)$. For $1 \leq k < i < j \leq n$, by (3.3) and the Octahedral Axiom we get the following commutative diagram with first two rows and the last two columns being distinguished triangles in A-$\underline{\text{mod}}$:

$$\begin{tikzcd}
\Omega Y_j^i \arrow[r] \arrow[d] & X_j \arrow[r, "\underline{\phi_i}\cdots\underline{\phi_{j-1}}"] \arrow[d, equal] & X_i \arrow[r] \arrow[d, "\underline{\phi_k}\cdots\underline{\phi_{i-1}}"] & Y_j^i \arrow[d] \\
\Omega Y_j^k \arrow[r] & X_j \arrow[r, "\underline{\phi_k}\cdots\underline{\phi_{j-1}}"] & X_k \arrow[r] \arrow[d] & Y_j^k \arrow[d] \\
 & & Y_i^k \arrow[r, equal] \arrow[d] & Y_i^k \arrow[d] \\
 & & \Omega^{-1}X_i \arrow[r] & \Omega^{-1}Y_j^i.
\end{tikzcd} \tag{3.11}$$

For $1 \leq m \leq n$ we prove the following formula by induction

$$\begin{aligned}\mathbf{Rot}^m X_{(\underline{\phi_i})} = \; & (\Omega^{-(m-2)}Y_m^{m-1} \to \Omega^{-(m-2)}Y_m^{m-2} \to \cdots \to \Omega^{-(m-2)}Y_m^1 \to \Omega^{-(m-1)}X_m \to \\ & \Omega^{-(m-1)}Y_n^m \to \Omega^{-(m-1)}Y_{n-1}^m \to \cdots \to \Omega^{-(m-1)}Y_{m+1}^m)\end{aligned} \tag{3.12}$$

Convention about (3.12): $\Omega^{-(m-1)}X_m$ is the $(n-m+1)$-st branch of $\mathbf{Rot}^m X_{(\underline{\phi_i})}$.

By definition (3.12) holds for $m = 1$. Assume that it holds for $1 \leq m \leq n-1$. Consider the following commutative diagram with rows of distinguished triangles

$$\begin{tikzcd}
\Omega^{-(m-2)}Y_m^{m-1} \arrow[r] \arrow[d] & \Omega^{-(m-1)}Y_{m+1}^m \arrow[r] \arrow[d, equal] & \Omega^{-(m-1)}Y_{m+1}^{m-1} \arrow[r] \arrow[d, dotted] & \Omega^{-(m-1)}Y_m^{m-1} \arrow[d] \\
\vdots \arrow[d] & \vdots \arrow[d, equal] & \vdots \arrow[d, dotted] & \vdots \arrow[d] \\
\Omega^{-(m-2)}Y_m^1 \arrow[r] \arrow[d] & \Omega^{-(m-1)}Y_{m+1}^m \arrow[r] \arrow[d, equal] & \Omega^{-(m-1)}Y_{m+1}^1 \arrow[r] \arrow[d, dotted] & \Omega^{-(m-1)}Y_m^1 \arrow[d] \\
\Omega^{-(m-1)}X_m \arrow[r] \arrow[d] & \Omega^{-(m-1)}Y_{m+1}^m \arrow[r] \arrow[d, equal] & \Omega^{-m}X_{m+1} \arrow[r] \arrow[d, dotted] & \Omega^{-m}X_m \arrow[d] \\
\Omega^{-(m-1)}Y_n^m \arrow[r] \arrow[d] & \Omega^{-(m-1)}Y_{m+1}^m \arrow[r] \arrow[d, equal] & \Omega^{-m}Y_n^{m+1} \arrow[r] \arrow[d, dotted] & \Omega^{-m}Y_n^m \arrow[d] \\
\vdots \arrow[d] & \vdots \arrow[d, equal] & \vdots \arrow[d, dotted] & \vdots \arrow[d] \\
\Omega^{-(m-1)}Y_{m+2}^m \arrow[r] & \Omega^{-(m-1)}Y_{m+1}^m \arrow[r] & \Omega^{-m}Y_{m+2}^{m+1} \arrow[r] & \Omega^{-m}Y_{m+2}^m
\end{tikzcd}$$

where the l-th row $(1 \le l \le m-1)$ is from the fourth column of (3.11) by taking $j = m+1, i = m, k = m-l$, and then applying $\Omega^{-(m-1)}$; the m-th row is from the first row of (3.11) by taking $j = m+1, i = m$, and then applying $\Omega^{-(m-1)}$; the l-th row $(m+1 \le l \le n-1)$ is from the fourth column of (3.11) by taking $j = n+m+1-l, i = m+1, k = m$, and then applying Ω^{-m}. By the definition of the rotation (cf. (3.4)), this proves (3.12) for $m+1$.

3.3. For $X_{(\underline{\phi_i})} \in \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$, define $\Omega^{-1}X_{(\underline{\phi_i})}$ to be $\begin{pmatrix}\Omega^{-1}X_1\\ \vdots \\ \Omega^{-1}X_n\end{pmatrix}_{(\Omega^{-1}\underline{\phi_i})} \in \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$.

Lemma 3.3. *Let* $X_{(\phi_i)} \in \mathrm{Mor}_n(A)$. *Then* $\mathbf{Rot}^{j(n+1)}X_{(\underline{\phi_i})} = \Omega^{-j(n-1)}X_{(\underline{\phi_i})}$, $\forall\ j \ge 1$.

Proof. By taking $m = n$ in (3.12), we get

$$\mathbf{Rot}^n X_{(\underline{\phi_i})} = (\Omega^{-(n-2)}Y_n^{n-1} \to \Omega^{-(n-2)}Y_n^{n-2} \to \cdots \to \Omega^{-(n-2)}Y_n^1 \to \Omega^{-(n-1)}X_n).$$

We have the following commutative diagram with rows being distinguished triangles

$$\begin{tikzcd}
\Omega^{-(n-2)}Y_n^{n-1} \arrow[r] \arrow[d] & \Omega^{-(n-1)}X_n \arrow[r, "(-1)^{n-1}\Omega^{-(n-1)}\underline{\phi_{n-1}}"] \arrow[d, equal] & \Omega^{-(n-1)}X_{n-1} \arrow[r] \arrow[d, dashed, "\Omega^{-(n-1)}\underline{\phi_{n-2}}"] & \Omega^{-(n-1)}Y_n^{n-1} \arrow[d] \\
\Omega^{-(n-2)}Y_n^{n-2} \arrow[r] \arrow[d] & \Omega^{-(n-1)}X_n \arrow[r, "(-1)^{n-1}\Omega^{-(n-1)}\underline{\phi_{n-2}}\ \underline{\phi_{n-1}}"] \arrow[d, equal] & \Omega^{-(n-1)}X_{n-2} \arrow[r] \arrow[d, dashed] & \Omega^{-(n-1)}Y_n^{n-2} \arrow[d] \\
\vdots \arrow[d] & \vdots \arrow[d, equal] & \vdots \arrow[d, dashed, "\Omega^{-(n-1)}\underline{\phi_2}"] & \vdots \arrow[d] \\
\Omega^{-(n-2)}Y_n^2 \arrow[r] \arrow[d] & \Omega^{-(n-1)}X_n \arrow[r, "(-1)^{n-1}\Omega^{-(n-1)}\underline{\phi_2}\cdots\underline{\phi_{n-1}}"] \arrow[d, equal] & \Omega^{-(n-1)}X_2 \arrow[r] \arrow[d, dashed, "\Omega^{-(n-1)}\underline{\phi_1}"] & \Omega^{-(n-1)}Y_n^2 \arrow[d] \\
\Omega^{-(n-2)}Y_n^1 \arrow[r] & \Omega^{-(n-1)}X_n \arrow[r, "(-1)^{n-1}\Omega^{-(n-1)}\underline{\phi_1}\cdots\underline{\phi_{n-1}}"] & \Omega^{-(n-1)}X_1 \arrow[r] & \Omega^{-(n-1)}Y_n^1
\end{tikzcd}$$

where the l-th row $(1 \le l \le n-1)$ is from the first row of (3.11) by taking $j = n, i = n-l$, and then applying $\Omega^{-(n-1)}$ (note that $(-1)^{(n-1)}$ arises from applying $\Omega^{-(n-1)}$). Using (3.4) we get that $\mathbf{Rot}^{n+1}X_{(\underline{\phi_i})}$ is

$$\begin{aligned}
&(\ \Omega^{-(n-1)}X_n \xrightarrow{(-1)^{n-1}\Omega^{-(n-1)}\underline{\phi_{n-1}}} \Omega^{-(n-1)}X_{n-1} \xrightarrow{\Omega^{-(n-1)}\underline{\phi_{n-2}}} \cdots \xrightarrow{\Omega^{-(n-1)}\underline{\phi_1}} \Omega^{-(n-1)}X_1\ )\\
&\cong (\ \Omega^{-(n-1)}X_n \xrightarrow{\Omega^{-(n-1)}\underline{\phi_{n-1}}} \Omega^{-(n-1)}X_{n-1} \xrightarrow{\Omega^{-(n-1)}\underline{\phi_{n-2}}} \cdots \xrightarrow{\Omega^{-(n-1)}\underline{\phi_1}} \Omega^{-(n-1)}X_1\ )\\
&= \Omega^{-(n-1)}X_{(\underline{\phi_i})}.
\end{aligned}$$

From this and induction the assertion follows. ■

3.4. Since τ is a triangle functor, by construction we see

$$\mathbf{Rot}\ \tau X_{(\underline{\phi_i})} \cong \tau\ \mathbf{Rot} X_{(\underline{\phi_i})}. \tag{3.13}$$

We have the following important result.

Theorem 3.4. *Let A be a selfinjective algebra, $X_{(\phi_i)} \in \mathcal{S}_n(A)$. Then there are the following isomorphisms in* $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$

(*i*) $\underline{\tau^j_{\mathcal{S}} X_{(\phi_i)}} \cong \tau^j\ \mathbf{Rot}^j X_{(\underline{\phi_i})}$ *for* $j \geq 1$. *In particular,* $\underline{\tau_{\mathcal{S}} X_{(\phi_i)}} \cong \tau\ \underline{\mathbf{Cok} X_{(\phi_i)}}$.

(*ii*) $\underline{\tau^{s(n+1)}_{\mathcal{S}} X_{(\phi_i)}} \cong \tau^{s(n+1)}\ \Omega^{-s(n-1)} X_{(\underline{\phi_i})},\ \forall\ s \geq 1.$

Proof. (*i*) First, we claim that there are the following isomorphisms in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$:

$$\underline{(\mathbf{Cok\ Mimo}\ \tau)^j\ Y_{(\psi_i)}} \cong \tau^j\ \mathbf{Rot}^j Y_{(\underline{\psi_i})}, \quad \forall\ Y_{(\psi_i)} \in \mathrm{Mor}_n(A\text{-mod}),\ \forall\ j \geq 1. \tag{3.14}$$

In fact, by Lemma 3.1 and induction we have

$$\begin{aligned}
\underline{(\mathbf{Cok\ Mimo}\ \tau)^j\ Y_{(\psi_i)}} &\cong \underline{\mathbf{Cok\ Mimo}\ \tau\ (\mathbf{Cok\ Mimo}\ \tau)^{j-1}\ Y_{(\psi_i)}} \\
&\overset{Lemma\ 3.1}{\cong} \mathbf{Rot}\ \underline{\tau\ (\mathbf{Cok\ Mimo}\ \tau)^{j-1}\ Y_{(\psi_i)}} \\
&\overset{(2.3)}{\cong} \mathbf{Rot}\ \tau\ \underline{(\mathbf{Cok\ Mimo}\ \tau)^{j-1}\ Y_{(\psi_i)}} \\
&\overset{Induction}{\cong} \mathbf{Rot}\ \tau^j\ \mathbf{Rot}^{j-1} Y_{(\underline{\psi_i})} \\
&\overset{(3.13)}{\cong} \tau^j\ \mathbf{Rot}^j Y_{(\underline{\psi_i})}.
\end{aligned}$$

Now, we have the following isomorphisms in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$:

$$\begin{aligned}
\underline{\tau^j_{\mathcal{S}} X_{(\phi_i)}} &\overset{Theorem\ 2.4}{\cong} \underline{(\mathbf{Mimo}\ \tau\ \mathbf{Cok})^j X_{(\phi_i)}} \\
&\cong \underline{\mathbf{Mimo}\ \tau\ (\mathbf{Cok\ Mimo}\ \tau)^{j-1}\ \mathbf{Cok} X_{(\phi_i)}} \\
&\overset{(2.3)}{\cong} \tau\ \underline{(\mathbf{Cok\ Mimo}\ \tau)^{j-1}\ \mathbf{Cok} X_{(\phi_i)}} \\
&\overset{(3.14)}{\cong} \tau^j\ \mathbf{Rot}^{j-1}\ \underline{\mathbf{Cok} X_{(\phi_i)}} \\
&\cong \tau^j\ \mathbf{Rot}^{j-1}\ \underline{\mathbf{Cok\ Mimo} X_{(\phi_i)}} \\
&\overset{Lemma\ 3.1}{\cong} \tau^j\ \mathbf{Rot}^j X_{(\underline{\phi_i})},
\end{aligned}$$

where we have used $\mathbf{Mimo} X_{(\phi_i)} = X_{(\phi_i)}$ since $X_{(\phi_i)} \in \mathcal{S}_n(A)$.

(*ii*) This follows from Lemma 3.3 and (*i*) by taking $j = s(n+1)$. ■

3.5. Now we pass from $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ to $\mathrm{Mor}_n(A\text{-mod})$. Before stating the main result, we need a notation. Let $X_{(\phi_i)} \in \mathrm{Mor}_n(A)$. For positive integers r and t, the object $\tau^r\ \Omega^{-t} X_{(\underline{\phi_i})} \in \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ is already defined (cf. 2.4 and 3.3). As in 2.4, we consider the full subcategory of $\mathrm{Mor}_n(A)$ given by

$$\{Y_{(\psi_i)} = \begin{pmatrix} \tau^r\ \Omega^{-t}X_1 \\ \vdots \\ \tau^r\ \Omega^{-t}X_n \end{pmatrix}_{(\psi_i)} \in \mathrm{Mor}_n(A) \mid Y_{(\underline{\psi_i})} \cong \tau^r\ \Omega^{-t}X_{(\underline{\phi_i})}\}.$$

Any object in this subcategory will be denoted by $\tau^r\ \Omega^{-t}X_{(\phi_i)}$ (we emphasize that this convention will cause no confusions). So, we have $\underline{\tau^r\ \Omega^{-t}X_{(\phi_i)}} \cong \tau^r\ \Omega^{-t}X_{(\underline{\phi_i})}$. By Lemma 1.6($i$), $\mathbf{Mimo}\ \tau^r\ \Omega^{-t}X_{(\phi_i)} \in \mathcal{S}_n(A)$ is a well-defined object, and there are the following isomorphisms in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$

$$\underline{\mathbf{Mimo}\ \tau^r\ \Omega^{-t}X_{(\phi_i)}} \cong \underline{\tau^r\ \Omega^{-t}X_{(\phi_i)}} \cong \tau^r\ \Omega^{-t}X_{(\underline{\phi_i})}. \tag{3.15}$$

Theorem 3.5. *Let A be a selfinjective algebra, and $X_{(\phi_i)} \in \mathcal{S}_n(A)$. Then we have*

$$\tau_{\mathcal{S}}^{s(n+1)}X_{(\phi_i)} \cong \mathbf{Mimo}\ \tau^{s(n+1)}\ \Omega^{-s(n-1)}X_{(\phi_i)}, \quad s \geq 1. \tag{3.16}$$

Proof. By Theorem 3.4(ii) we have

$$\underline{\tau_{\mathcal{S}}^{s(n+1)}X_{(\phi_i)}} \cong \tau^{s(n+1)}\ \Omega^{-s(n-1)}X_{(\underline{\phi_i})} \cong \underline{\mathbf{Mimo}\ \tau^{s(n+1)}\Omega^{-s(n-1)}X_{(\phi_i)}}.$$

Since $\mathbf{Mimo}\ \tau^{s(n+1)}\ \Omega^{-s(n-1)}X_{(\phi_i)} \in \mathcal{S}_n(A)_{\mathcal{I}}$ (cf. Remark (ii) in 1.4), the assertion follows from Corollary 2.6. ■

3.6. We apply Theorem 3.4 to the algebra $\Lambda(m,t)$, which is defined below. Let $\mathbb{Z}_m$ be the cyclic quiver with vertices indexed by the cyclic group $\mathbb{Z}/m\mathbb{Z}$ of order m, and with arrows $a_i:\ i \longrightarrow i+1,\ \forall\ i \in \mathbb{Z}/m\mathbb{Z}$. Let $k\mathbb{Z}_m$ be the path algebra of the quiver $\mathbb{Z}_m$, J the ideal generated by all arrows, and $\Lambda(m,t) := k\mathbb{Z}_m/J^t$ with $m \geq 1,\ t \geq 2$. Any connected selfinjective Nakayama algebra over an algebraically closed field is Morita equivalent to $\Lambda(m,t)$, $m \geq 1,\ t \geq 2$. Note that $\Lambda(m,t)$ is a Frobenius algebra of finite representation type, and that $\Lambda(m,t)$ is symmetric if and only if $m \mid (t-1)$. For the Auslander-Reiten sequences of $\Lambda(m,t)$ see [ARS], p.197. In the stable category $\Lambda(m,t)$-$\underline{\mathrm{mod}}$, we have the following information on the orders of τ and Ω (see 5.1 in [CZ])

$$o(\tau) = m; \qquad o(\Omega) = \begin{cases} m, & t = 2; \\ \frac{2m}{(m,t)}, & t \geq 3, \end{cases} \tag{3.17}$$

where (m,t) is the g.c.d of m and t. By (3.16) and (3.17) we get the following

Corollary 3.6. *For an indecomposable nonprojective object $X_{(\phi_i)} \in \mathcal{S}_n(\Lambda(m,t))$, $m \geq 1,\ t \geq 2$, there are the following isomorphisms:*

(i) *If n is odd, then $\tau_{\mathcal{S}}^{m(n+1)}X_{(\phi_i)} \cong X_{(\phi_i)}$;*

(ii) *If n is even, then $\tau_{\mathcal{S}}^{2m(n+1)}X_{(\phi_i)} \cong X_{(\phi_i)}$.*

3.7. **Example.** Let $A = kQ/\langle \delta\alpha, \beta\gamma, \alpha\delta - \gamma\beta\rangle$, where Q is the quiver $2\bullet \underset{\delta}{\overset{\alpha}{\rightleftarrows}} 1\bullet \underset{\gamma}{\overset{\beta}{\rightleftarrows}} 3\bullet$

Then A is selfinjective with $\tau \cong \Omega^{-1}$ and $\Omega^6 \cong \mathrm{id}$ on the object of A-$\underline{\mathrm{mod}}$. The Auslander-Reiten quiver of A is



Let $X_{(\phi_i)}$ be an indecomposable nonprojective object in $\mathcal{S}_n(A)$. By (3.16), for $s \ge 1$ we have $\tau_{\mathcal{S}}^{s(n+1)} X_{(\phi_i)} \cong \textbf{Mimo}\ \tau^{s(n+1)}\ \Omega^{-s(n-1)} X_{(\phi_i)} \cong \textbf{Mimo}\ \Omega^{-2sn} X_{(\phi_i)}$ in $\mathcal{S}_n(A)$. Then by Remark (i) in 1.4 we get

(i) if $n \equiv 0,$ or 3 (mod6), then $\tau_{\mathcal{S}}^{n+1} X_{(\phi_i)} \cong X_{(\phi_i)}$; and

(ii) if $n \equiv \pm 1,$ or ± 2 (mod6), then $\tau_{\mathcal{S}}^{3(n+1)} X_{(\phi_i)} \cong X_{(\phi_i)}$.

4. **Serre functors of stable monomorphism categories**

Throughout this section, A is a finite-dimensional selfinjective algebra over a field. We study the periodicity of the Serre functor $F_{\mathcal{S}}$ on the objects of the stable monomorphism category $\underline{\mathcal{S}_n(A)}$. In particular, $F_{\mathcal{S}}^{m(n+1)} X \cong X$ for $X \in \underline{\mathcal{S}_n(\Lambda(m,t))}$.

4.1. Let $\mathcal{A}$ be a Hom-finite Krull-Schmidt triangulated k-category with suspension functor [1]. For the Auslander-Reiten triangles we refer to [H]. In an Auslander-Reiten triangle $X \to Y \to Z \to X[1]$, the indecomposable object X is uniquely determined by Z. Write $X = \widetilde{\tau_{\mathcal{A}}} Z$, and extend the action of $\widetilde{\tau_{\mathcal{A}}}$ to arbitrary objects, and put $\widetilde{\tau_{\mathcal{A}}} 0 = 0$. In general, $\widetilde{\tau_{\mathcal{A}}}$ is *not* a functor. By Theorem I.2.4 of [RV], $\mathcal{A}$ has a Serre functor F if and if $\mathcal{A}$ has Auslander-Reiten triangles; if this is the case, F and $[1]\widetilde{\tau_{\mathcal{A}}}$ coincide on the objects of $\mathcal{A}$. If $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} X[1]$ is an Auslander-Reiten triangle, then so is $X[1] \xrightarrow{-f[1]} Y[1] \xrightarrow{-g[1]} Z[1] \xrightarrow{-h[1]} X[2]$. It follows that

$$FZ \cong ([1]\widetilde{\tau_{\mathcal{A}}})Z \cong (\widetilde{\tau_{\mathcal{A}}}[1])Z, \ \forall\ Z \in \mathcal{A}. \tag{4.1}$$

4.2. Since A is self-injective, by Corollary 4.1(ii) of [Z], $\mathcal{S}_n(A)$ is exactly the category of Gorenstein projective $T_n(A)$-modules, hence it is a Frobenius category whose projective-injective objects are exactly all the projective $T_n(A)$-modules. Thus, the stable category $\underline{\mathcal{S}_n(A)}$ of $\mathcal{S}_n(A)$ modulo projective objects is a Hom-finite Krull-Schmidt triangulated category with suspension functor $\Omega_{\mathcal{S}}^{-1} = \Omega_{\mathcal{S}_n(A)}^{-1}$. Since $\mathcal{S}_n(A)$ has Auslander-Reiten sequences, it follows that $\underline{\mathcal{S}_n(A)}$ has Auslander-Reiten triangles, and hence it has a Serre functor $F_{\mathcal{S}} = F_{\mathcal{S}_n(A)}$, which coincides with $\Omega_{\mathcal{S}}^{-1}\widetilde{\tau_{\mathcal{S}}}$ on the objects of $\underline{\mathcal{S}_n(A)}$.

In order to make the following computation more clear, we denote by $Q : \mathcal{S}_n(A) \rightarrow \underline{\mathcal{S}_n(A)}$ the natural functor. Then

$$\widetilde{\tau_{\mathcal{S}}}\ QZ = Q\ \tau_{\mathcal{S}}Z = \tau_{\mathcal{S}}Z,\ \forall Z \in \mathcal{S}_n(A). \tag{4.2}$$

4.3. In 2.6 we have considered the canonical functor $W : \mathcal{S}_n(A)_{\mathcal{I}} \rightarrow \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ given by $X_{(\phi_i)} \mapsto X_{(\underline{\phi_i})}$.

Lemma 4.1. *The functor W induces a functor $\widetilde{W} : \underline{\mathcal{S}_n(A)} \rightarrow \mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ satisfying $\widetilde{W}\ Q|_{\mathcal{S}_n(A)_{\mathcal{I}}} = W$; and $\widetilde{W}$ reflects isomorphisms.*

Proof. The definition of $\widetilde{W}$ is clear by the requirement $\widetilde{W}Q|_{\mathcal{S}_n(A)_{\mathcal{I}}} = W$. We need to check that it is well-defined. If $X_{(\phi_i)}$ and $Y_{(\psi_i)}$ are indecomposable and nonprojective in $\mathcal{S}_n(A)$, and $X_{(\phi_i)} \cong Y_{(\psi_i)}$ in $\underline{\mathcal{S}_n(A)}$, then $X_{(\phi_i)} \cong Y_{(\psi_i)}$ in $\mathcal{S}_n(A)$, and hence $X_{(\underline{\phi_i})} \cong Y_{(\underline{\psi_i})}$ in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$, i.e., $\widetilde{W}$ is well-defined on objects. For any morphism $f = (f_i) : X_{(\phi_i)} \rightarrow Y_{(\psi_i)}$ in $\mathcal{S}_n(A)$ which factors through a projective object of $\mathcal{S}_n(A)$, by Lemma 1.1(iii), the morphism $(\underline{f_i}) = 0$ in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$. Thus $\widetilde{W}$ is well-defined.

Assume that $\widetilde{W}X \cong \widetilde{W}Y$ in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ for $X, Y \in \underline{\mathcal{S}_n(A)}$. We may write $X = QX'$, $Y = QY'$ with $X', Y' \in \mathcal{S}_n(A)_{\mathcal{I}}$. Then $WX' \cong WY'$ in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$. By Corollary 2.6, W reflects isomorphisms, thus $X' \cong Y'$ in $\mathcal{S}_n(A)_{\mathcal{I}}$, and hence $X \cong Y$ in $\underline{\mathcal{S}_n(A)}$. ■

Lemma 4.2. *For $X = X_{(\phi_i)} \in \underline{\mathcal{S}_n(A)}$, we have the following isomorphism in* $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$

$$\widetilde{W}\ \Omega_{\mathcal{S}}X \cong \Omega\ \widetilde{W}X. \tag{4.3}$$

Proof. Let $0 \rightarrow \Omega_{\mathcal{S}}X \rightarrow P \rightarrow X \rightarrow 0$ be an exact sequence in $\mathcal{S}_n(A)$ with P projective. Taking the i-th branches we see that $(\Omega_{\mathcal{S}}X)_i = \Omega X_i \oplus P_i'$ for some projective A-module P_i'. It follows that $(\Omega_{\mathcal{S}}X)_i = \Omega X_i$ in A-$\underline{\mathrm{mod}}$. Write $\Omega_{\mathcal{S}}X$ as $(\Omega_{\mathcal{S}}X)_{(\psi_i)}$. By the following commutative diagram with exact columns

$$\begin{tikzcd}
0 \arrow[d] & 0 \arrow[d] & & 0 \arrow[d] & 0 \arrow[d] \\
(\Omega_{\mathcal{S}}X)_n \arrow[r, "\psi_{n-1}"] \arrow[d] & (\Omega_{\mathcal{S}}X)_{n-1} \arrow[r] \arrow[d] & \cdots \arrow[r] & (\Omega_{\mathcal{S}}X)_2 \arrow[r, "\psi_1"] \arrow[d] & (\Omega_{\mathcal{S}}X)_1 \arrow[d] \\
P_n \arrow[r] \arrow[d] & P_{n-1} \arrow[r] \arrow[d] & \cdots \arrow[r] & P_2 \arrow[r] \arrow[d] & P_1 \arrow[d] \\
X_n \arrow[r, "\phi_{n-1}"] \arrow[d] & X_{n-1} \arrow[r] \arrow[d] & \cdots \arrow[r] & X_2 \arrow[r, "\phi_1"] \arrow[d] & X_1 \arrow[d] \\
0 & 0 & & 0 & 0
\end{tikzcd}$$

we see $\underline{\psi_i} = \Omega\underline{\phi_i}$, and hence the assertion follows. ■

4.4. For positive integers a and b, let $[a,b]$ denote the l.c.m of a and b. The main result of this section is

Theorem 4.3. *Let A be a selfinjective algebra, and $F_{\mathcal{S}}$ be the Serre functor of $\underline{\mathcal{S}_n(A)}$. Then we have an isomorphism in $\underline{\mathcal{S}_n(A)}$ for $X_{(\phi_i)} \in \underline{\mathcal{S}_n(A)}$ and for $s \geq 1$*

$$F_{\mathcal{S}}^{s(n+1)} X_{(\phi_i)} \cong \mathbf{Mimo}\ \tau^{s(n+1)}\ \Omega^{-2sn} X_{(\phi_i)}. \tag{4.4}$$

Moreover, if d_1 and d_2 are positive integers such that $\tau^{d_1} M \cong M$ and $\Omega^{d_2} M \cong M$ for each indecomposable nonprojective A-module M, then $F_{\mathcal{S}}^{N(n+1)} X_{(\phi_i)} \cong X_{(\phi_i)}$, where $N = [\frac{d_1}{(n+1,d_1)}, \frac{d_2}{(2n,d_2)}]$.

Proof. We have isomorphisms in $\mathrm{Mor}_n(A\text{-}\underline{\mathrm{mod}})$ for $s \geq 1$:

$$\begin{aligned}
\widetilde{W}\ F_{\mathcal{S}}^{s(n+1)} X_{(\phi_i)} &\overset{(4.1)}{\cong} \widetilde{W}\ \Omega_{\mathcal{S}}^{-s(n+1)}\ \widetilde{\tau_{\mathcal{S}}}^{s(n+1)} X_{(\phi_i)} \\
&\overset{(4.3)}{\cong} \Omega^{-s(n+1)}\ \widetilde{W}\ \widetilde{\tau_{\mathcal{S}}}^{s(n+1)} X_{(\phi_i)} \\
&\overset{(4.2)}{\cong} \Omega^{-s(n+1)}\ \widetilde{W}\ Q\ \tau_{\mathcal{S}}^{s(n+1)} X_{(\phi_i)} \\
&\overset{(3.16)}{\cong} \Omega^{-s(n+1)}\ \widetilde{W}\ Q\ \mathbf{Mimo}\ \tau^{s(n+1)}\ \Omega^{-s(n-1)} X_{(\phi_i)} \\
&\overset{Lemma\ 4.1}{\cong} \Omega^{-s(n+1)}\ W\ \mathbf{Mimo}\ \tau^{s(n+1)}\ \Omega^{-s(n-1)} X_{(\phi_i)} \\
&\cong \Omega^{-s(n+1)}\ \underline{\tau^{s(n+1)}\ \Omega^{-s(n-1)} X_{(\phi_i)}} \\
&\overset{(3.15)}{\cong} \Omega^{-s(n+1)}\ \tau^{s(n+1)}\ \Omega^{-s(n-1)} X_{(\underline{\phi_i})} \\
&\cong \tau^{s(n+1)}\ \Omega^{-2sn} X_{(\underline{\phi_i})} \\
&\overset{(3.15)}{\cong} \widetilde{W}\ \mathbf{Mimo}\ \tau^{s(n+1)}\ \Omega^{-2sn} X_{(\phi_i)}.
\end{aligned}$$

Now (4.4) follows from Lemma 4.1. Since $d_1 | N(n+1)$, $d_2 | (2Nn)$, taking $s = N$ in (4.4) we get $F_{\mathcal{S}}^{N(n+1)} X_{(\phi_i)} \cong \mathbf{Mimo}\ \tau^{N(n+1)}\ \Omega^{-2Nn} X_{(\phi_i)} \cong \mathbf{Mimo} X_{(\phi_i)} = X_{(\phi_i)}$. ■

Note that the conditions on τ and Ω in Theorem 4.3 hold in particular for representation-finite selfinjective algebras.

4.5. Applying Theorem 4.3 to the selfinjective Nakayama algebras $\Lambda(m,t)$, we get

Corollary 4.4. *Let $F_{\mathcal{S}}$ be the Serre functor of $\underline{\mathcal{S}_n(\Lambda(m,t))}$ with $m \geq 1,\ t \geq 2$, and X be an arbitrary object in $\underline{\mathcal{S}_n(\Lambda(m,t))}$. Then*

(i) If $t = 2$, then $F_{\mathcal{S}}^{N(n+1)} X \cong X$, where $N = \frac{m}{(m,n-1)}$.

(ii) If $t \geq 3$, then $F_{\mathcal{S}}^{N(n+1)} X \cong X$, where $N = \frac{m}{(m,t,n+1)}$.

Proof. (i) In this case $\tau = \Omega$, and $o(\tau) = m = o(\Omega)$, by (3.17). Put $N = \frac{m}{(m,n-1)}$. It follows from (4.4) that

$$F_{\mathcal{S}}^{N(n+1)}X \cong \mathbf{Mimo}\ \tau^{N(n+1)}\ \Omega^{-2Nn}X \cong \mathbf{Mimo}\ \Omega^{-N(n-1)}X$$

$$\cong \mathbf{Mimo}\ \Omega^{-m\frac{n-1}{(m,n-1)}}X \cong \mathbf{Mimo}X = X.$$

(ii) This follows from Theorem 4.3 by taking $d_1 = m$ and $d_2 = \frac{2m}{(m,t)}$. By a computation in elementary number theory, we get

$$[\frac{d_1}{(n+1,d_1)}, \frac{d_2}{(2n,d_2)}] = [\frac{m}{(n+1,m)}, \frac{\frac{2m}{(m,t)}}{(2n, \frac{2m}{(m,t)})}] = \frac{m}{(m,t,n+1)}. \qquad \blacksquare$$

5. Appendix 1: Proofs of Lemmas 1.5 and 1.6

We give proofs of Lemmas 1.5 and 1.6.

Lemma 5.1. *Let* $X_{(\phi_i)} \in \mathrm{Mor}_n(A)$, $I_2, \cdots, I_n$ *be injective* A*-modules such that* $X'_{(\phi'_i)} = \begin{pmatrix} X_1\oplus I_2\oplus\cdots\oplus I_n \\ \vdots \\ X_{n-1}\oplus I_n \\ X_n \end{pmatrix}_{(\phi'_i)} \in \mathcal{S}_n(A)$, *where* $\phi'_i = \begin{pmatrix} \phi_i & 0 & 0 & \cdots & 0 \\ * & 0 & 0 & \cdots & 0 \\ * & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ * & 0 & 0 & \cdots & 1 \end{pmatrix}_{(n-i+1)\times(n-i)}$. *Then* $X'_{(\phi'_i)} \cong \mathbf{Mimo}X_{(\phi_i)} \oplus J$, *where* J *is an injective object of* $\mathcal{S}_n(A)$. *Moreover,* $J = \bigoplus\limits_{i=1}^{n-1} \mathbf{m}_i(Q_i)$, *where* Q_i *is an injective* A*-module such that* $Q_i \oplus \mathrm{IKer}\phi_i \cong I_{i+1}$, $1 \le i \le n-1$.

Proof. It is clear that the morphism $\begin{pmatrix} (1,0,\cdots,0) \\ \vdots \\ (1,0) \\ 1 \end{pmatrix} : X'_{(\phi'_i)} \twoheadrightarrow X_{(\phi_i)}$ is a right approximation of $X_{(\phi_i)}$ in $\mathcal{S}_n(A)$ (this can be proved as Lemma 1.3(i), see [Z], Lemma 2.3). By Lemma 1.3(i), there is an object $J \in \mathcal{S}_n(A)$ such that $X'_{(\phi'_i)} \cong \mathbf{Mimo}X_{(\phi_i)} \oplus J$. Comparing the branches we get $J_n = 0$ and

$$I_{i+1} \oplus \cdots \oplus I_n \cong \mathrm{IKer}\phi_i \oplus \cdots \oplus \mathrm{IKer}\phi_{n-1} \oplus J_i, \ \forall\ 1 \le i \le n-1. \tag{5.1}$$

Put $Q_{n-1} = J_{n-1}$. From $I_n \cong \mathrm{IKer}\phi_{n-1} \oplus J_{n-1}$ we see that Q_{n-1} is an injective A-module. Since $J \in \mathcal{S}_n(A)$, Q_{n-1} is a submodule of J_{n-2}, thus $J_{n-2} = Q_{n-2} \oplus Q_{n-1}$. By $I_{n-1} \oplus I_n \cong \mathrm{IKer}\phi_{n-2} \oplus \mathrm{IKer}\phi_{n-1} \oplus J_{n-2}$ in (5.1), we see $I_{n-1} \cong \mathrm{IKer}\phi_{n-2} \oplus Q_{n-2}$. Repeating this process we see that J_i is of the form $J_i = Q_i \oplus \cdots \oplus Q_{n-1}$ with Q_i being injective A-modules, and $Q_i \oplus \mathrm{IKer}\phi_i \cong I_{i+1}$, $1 \le i \le n-1$. Thus $J = \bigoplus\limits_{i=1}^{n-1} \mathbf{m}_i(Q_i)$ is an injective object of $\mathcal{S}_n(A)$. $\blacksquare$

Now we can prove Lemma 1.5 (cf. Claim 2 in §4 of [RS2] for the case of $n = 2$).

Proof of Lemma 1.5. We just prove (i). Since the A-map $\phi'_i : X_{i+1} \oplus I_{i+2} \oplus \cdots \oplus I_n \to X_i \oplus I_{i+1} \oplus I_{i+2} \oplus \cdots \oplus I_n$ is monic, the restriction of ϕ'_i on $I_{i+2} \oplus \cdots \oplus I_n$ is also monic, and hence it is a split monomorphism. Hence $X'_{(\phi'_i)}$ is isomorphic to

$$X''_{(\phi''_i)} = \begin{pmatrix} X_1\oplus I_2\oplus\cdots\oplus I_n \\ \vdots \\ X_{n-1}\oplus I_n \\ X_n \end{pmatrix}_{(\phi''_i)} \in \mathcal{S}_n(A), \quad \text{where} \quad \phi''_i = \begin{pmatrix} \phi_i & 0 & 0 & \cdots & 0 \\ * & 0 & 0 & \cdots & 0 \\ * & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ * & 0 & 0 & \cdots & 1 \end{pmatrix}_{(n-i+1)\times(n-i)} .$$

Then the assertion follows from Lemma 5.1. ■

Lemma 5.2. *Let $X_{(f_i)}$, $X_{(g_i)} \in \mathrm{Mor}_n(A)$ such that $f_i - g_i$ factors through an injective A-module, $1 \le i \le n-1$, and $h_i : X_i \to I_i$ be an injective envelope, $2 \le i \le n$. Set*

$$X'_{(f'_i)} = \begin{pmatrix} X_1\oplus I_2\oplus\cdots\oplus I_n \\ \vdots \\ X_{n-1}\oplus I_n \\ X_n \end{pmatrix}_{(f'_i)} \quad \text{where} \quad f'_i = \begin{pmatrix} f_i & 0 & 0 & \cdots & 0 \\ h_{i+1} & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}_{(n-i+1)\times(n-i)} ,$$

and

$$X'_{(g'_i)} = \begin{pmatrix} X_1\oplus I_2\oplus\cdots\oplus I_n \\ \vdots \\ X_{n-1}\oplus I_n \\ X_n \end{pmatrix}_{(g'_i)} \quad \text{where} \quad g'_i = \begin{pmatrix} g_i & 0 & 0 & \cdots & 0 \\ h_{i+1} & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}_{(n-i+1)\times(n-i)} .$$

Then $X'_{(f'_i)} \cong X'_{(g'_i)}$ in $\mathcal{S}_n(A)$.

Proof. For $1 \le i \le n-1$, it is clear that $f_i - g_i : X_{i+1} \to X_i$ factors through the injective envelope $h_{i+1} : X_{i+1} \to I_{i+1}$, and hence there is an A-map $u_i : I_{i+1} \to X_i$ such that $g_i - f_i = u_i h_{i+1}$. The following commutative diagram shows $X'_{(f'_i)} \cong X'_{(g'_i)}$.

$$\begin{tikzcd}
X_n \arrow[r, "f'_{n-1}"] \arrow[d, equal, "\alpha_n = 1_{X_n}"'] & X_{n-1}\oplus I_n \arrow[r] \arrow[d, "\alpha_{n-1}"] & \cdots \arrow[r] & X_2\oplus I_3\oplus\cdots\oplus I_n \arrow[r, "f'_1"] \arrow[d, "\alpha_2"] & X_1\oplus I_2\oplus\cdots\oplus I_n \arrow[d, "\alpha_1"] \\
X_n \arrow[r, "g'_{n-1}"] & X_{n-1}\oplus I_n \arrow[r] & \cdots \arrow[r] & X_2\oplus I_3\oplus\cdots\oplus I_n \arrow[r, "g'_1"] & X_1\oplus I_2\oplus\cdots\oplus I_n
\end{tikzcd}$$

$$\text{where } \alpha_i = \begin{pmatrix} 1 & u_i & g_i u_{i+1} & \cdots & (g_i g_{i+1} \cdots g_{n-2} u_{n-1}) \\ 0 & 1 & h_{i+1} u_{i+1} & \cdots & (h_{i+1} g_{i+1} \cdots g_{n-2} u_{n-1}) \\ 0 & 0 & 1 & \cdots & (h_{i+2} g_{i+2} \cdots g_{n-2} u_{n-1}) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & h_{n-1} u_{n-1} \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}_{(n-i+1)\times(n-i+1)} , \ 1 \le i \le n.$$

■

Lemma 5.3. *Let $X_{(f_i)}$, $X_{(g_i)} \in \mathrm{Mor}_n(A)$ such that $f_i - g_i$ factors through an injective A-module, $1 \le i \le n-1$. If each X_i has no nonzero injective direct summands, then* $\mathbf{Mimo}X_{(f_i)} \cong \mathbf{Mimo}X_{(g_i)}$ *in $\mathcal{S}_n(A)$.*

Proof. Consider $X'_{(f'_i)}$ and $X'_{(g'_i)}$ defined in Lemma 5.2, which are isomorphic in $\mathcal{S}_n(A)$. By Lemma 5.1, there exist injective A-modules $Q_{f,i}$ and $Q_{g,i}$ such that $\mathrm{IKer} f_i \oplus Q_{f,i} \cong I_{i+1} \cong \mathrm{IKer} g_i \oplus Q_{g,i}$, $1 \le i \le n-1$, and

$$\mathbf{Mimo}X_{(f_i)} \oplus \bigoplus_{i=1}^{n-1} \mathbf{m}_i(Q_{f,i}) \cong X'_{(f'_i)} \cong X'_{(g'_i)} \cong \mathbf{Mimo}X_{(g_i)} \oplus \bigoplus_{i=1}^{n-1} \mathbf{m}_i(Q_{g,i}).$$

By Claim 3 in §4 of [RS2], we have $\operatorname{IKer} f_i \cong \operatorname{IKer} g_i,\ 1 \le i \le n-1$. Thus $Q_{f,i} \cong Q_{g,i}$, and $\bigoplus\limits_{i=1}^{n-1} \mathbf{m}_i(Q_{f,i}) \cong \bigoplus\limits_{i=1}^{n-1} \mathbf{m}_i(Q_{g,i})$, from which the assertion follows. ■

Now we can prove Lemma 1.6 (cf. Theorem 4.2 of [RS2] for the case of $n = 2$).

Proof of Lemma 1.6. We only prove (i). If $\mathbf{Mimo}X_{(\phi_i)} \cong \mathbf{Mimo}Y_{(\psi_i)}$ in $\mathcal{S}_n(A)$, then by the construction of **Mimo**, $X_{\overline{(\phi_i)}} \cong Y_{\overline{(\psi_i)}}$ in $\operatorname{Mor}_n(A\text{-}\overline{\text{mod}})$. Conversely, assume that $X_{\overline{(\phi_i)}} \cong Y_{\overline{(\psi_i)}}$ in $\operatorname{Mor}_n(A\text{-}\overline{\text{mod}})$. Since all X_i and Y_i have no nonzero injective direct summands, there are A-isomorphisms $x_i : X_i \to Y_i$, such that each $x_i\phi_i - \psi_i x_{i+1}$ factors through an injective A-module, $1 \le i \le n-1$. Then each $\phi_i - x_i^{-1}\psi_i x_{i+1}$ factors through an injective A-module, $1 \le i \le n-1$. By Lemma 5.3, $\mathbf{Mimo}X_{(x_i^{-1}\psi_i x_{i+1})} \cong \mathbf{Mimo}X_{(\phi_i)}$. Since $Y_{(\psi_i)} \cong X_{(x_i^{-1}\psi_i x_{i+1})}$, we get $\mathbf{Mimo}Y_{(\psi_i)} \cong \mathbf{Mimo}X_{(\phi_i)}$. ■

6. Appendix 2: Auslander-Reiten quivers of some monomorphism categories

We include the Auslander-Reiten quivers of some representation-finite monomorphism categories.

6.1. By Simson [S], $\mathcal{S}_{n,2}$, $\mathcal{S}_{2,3}$, $\mathcal{S}_{2,4}$, $\mathcal{S}_{2,5}$, $\mathcal{S}_{3,3}$ and $\mathcal{S}_{4,3}$ are the only representation-finite cases among all $\mathcal{S}_{n,t} = \mathcal{S}_n(k[x]/\langle x^t\rangle),\ n \ge 2,\ t \ge 2$. In [RS3], Ringel and Schmidmeier give the Auslander-Reiten quivers of $\mathcal{S}_{2,t},\ t = 2, 3, 4, 5$. In the following we give the remaining cases, namely, $\mathcal{S}_{n,2}\ \ (n \ge 3)$, $\ \mathcal{S}_{3,3}$, and $\mathcal{S}_{4,3}$.

(i) There are n indecomposable projective objects and $\frac{n(n+1)}{2}$ indecomposable nonprojective objects in $\mathcal{S}_{n,2}$. For the Auslander-Reiten quivers of $\mathcal{S}_{3,2}$ see 2.5. The Auslander-Reiten quiver of $\mathcal{S}_{4,2}$ is as follows, where $A = k[x]/\langle x^2\rangle$ and S is the simple A-module.



(ii) Let $A = k[x]/\langle x^3\rangle$. Denote by M and S the two indecomposable nonprojective A-modules, where S is simple. There are 3 indecomposable projective objects and 24 indecomposable nonprojective objects in $\mathcal{S}_{3,3}$, whose Auslander-Reiten quiver is as follows.



(*iii*) Let A, M, S be as in (*ii*). There are 4 indecomposable projective objects and 80 indecomposable nonprojective objects in $\mathcal{S}_{4,3}$, whose Auslander-Reiten quiver looks like.



6.2. Consider the selfinjective Nakayama algebra $\Lambda = \Lambda(2,2)$, whose indecomposable modules are denoted by $S_1 = \ k \rightleftarrows 0$, $S_2 = \ 0 \rightleftarrows k$, $P_1 = \ k \underset{0}{\overset{1}{\rightleftarrows}} k$, $P_2 = \ k \underset{1}{\overset{0}{\rightleftarrows}} k.$

There are 4 indecomposable projective objects and 6 indecomposable nonprojective objects in $\mathcal{S}_2(\Lambda)$, with the Auslander-Reiten quiver as follows.

$$\begin{tikzcd}
& \begin{smallmatrix}P_2\\0\end{smallmatrix} \arrow[dr] & & \begin{smallmatrix}P_2\\P_2\end{smallmatrix} \arrow[dr] & & & & \\
\begin{smallmatrix}S_1\\0\end{smallmatrix} \arrow[ur] \arrow[dr] & & \begin{smallmatrix}P_2\\S_1\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}S_2\\S_2\end{smallmatrix} \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}S_1\\0\end{smallmatrix} \arrow[dr] \arrow[ll, dotted] & \\
& \begin{smallmatrix}S_1\\S_1\end{smallmatrix} \arrow[ur] & & \begin{smallmatrix}S_2\\0\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}P_1\\S_2\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}S_1\\S_1\end{smallmatrix} \arrow[ll, dotted] \\
& & & & \begin{smallmatrix}P_1\\0\end{smallmatrix} \arrow[ur] & & \begin{smallmatrix}P_1\\P_1\end{smallmatrix} \arrow[ur] &
\end{tikzcd}$$

There are 6 indecomposable projective objects and 12 indecomposable nonprojective objects in $\mathcal{S}_3(\Lambda)$, with the Auslander-Reiten quiver as follows.

$$\begin{tikzcd}
& \begin{smallmatrix}P_2\\0\\0\end{smallmatrix} \arrow[dr] & & \begin{smallmatrix}P_2\\P_2\\0\end{smallmatrix} \arrow[dr] & & \begin{smallmatrix}P_2\\P_2\\P_2\end{smallmatrix} \arrow[dr] & & & & & \\
\begin{smallmatrix}S_1\\0\\0\end{smallmatrix} \arrow[ur] \arrow[dr] & & \begin{smallmatrix}P_2\\S_1\\0\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}P_2\\P_2\\S_1\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}S_2\\S_2\\S_2\end{smallmatrix} \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}S_1\\0\\0\end{smallmatrix} \arrow[dr] \arrow[ll, dotted] & & \\
& \begin{smallmatrix}S_1\\S_1\\0\end{smallmatrix} \arrow[ur] \arrow[dr] & & \begin{smallmatrix}P_2\\S_1\\S_1\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}S_2\\S_2\\0\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}P_1\\S_2\\S_2\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}S_1\\S_1\\0\end{smallmatrix} \arrow[dr] \arrow[ll, dotted] & \\
& & \begin{smallmatrix}S_1\\S_1\\S_1\end{smallmatrix} \arrow[ur] & & \begin{smallmatrix}S_2\\0\\0\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}P_1\\S_2\\0\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}P_1\\P_1\\S_2\end{smallmatrix} \arrow[ur] \arrow[dr] \arrow[ll, dotted] & & \begin{smallmatrix}S_1\\S_1\\S_1\end{smallmatrix} \arrow[ll, dotted] \\
& & & & & \begin{smallmatrix}P_1\\0\\0\end{smallmatrix} \arrow[ur] & & \begin{smallmatrix}P_1\\P_1\\0\end{smallmatrix} \arrow[ur] & & \begin{smallmatrix}P_1\\P_1\\P_1\end{smallmatrix} \arrow[ur] &
\end{tikzcd}$$

References

[Ar] D. M. Arnold, Abelian groups and representations of finite partially ordered sets, Canad. Math. Soc. Books in Math., Springer-Verlag, New York, 2000.

[ARS] M. Auslander, I. Reiten, S. O. Smalø, Representation theory of Artin algebras, Cambridge Studies in Adv. Math. 36., Cambridge Univ. Press, 1995.

[AS] M. Auslander, S. O. Smalø, Almost split sequences in subcategories, J. Algebra 69(1981), 426-454.

[B] G. Birkhoff, Subgroups of abelian groups, Proc. Lond. Math. Soc. II, Ser. 38(1934), 385-401.

[C] X. W. Chen, Stable monomorphism category of Frobenius category, avaible in arXiv: math. RT 0911.1987, 2009.

[CZ] C. Cibils, P.Zhang, Calabi-Yau objects in triangulated categories, Trans. Amer. Math. Soc. 361(2009), 6501-6519.

[H] D. Happel, Triangulated categories in representation theory of finite dimensional algebras, Lond. Math. Soc. Lecture Notes Ser. 119, Cambridge Univ. Press, 1988.

[KLM] D. Kussin, H. Lenzing, H. Meltzer, Nilpotent operators and weighted projective lines, avaible in arXiv: math. RT 1002.3797.

[RV] I. Reiten, M. Van den Bergh, Noetherian hereditary abelian categories satisfying Serre duality, J. Amer. Math. Soc. 15(2)(2002), 295-366.

[RW] F. Richman, E. A. Walker, Subgroups of p^5-bounded groups, in: Abelian groups and modules, Trends Math., Birkhäuser, Basel, 1999, 55-73.

[RS1] C. M. Ringel, M. Schmidmeier, Submodules categories of wild representation type, J. Pure Appl. Algebra 205(2)(2006), 412-422.

[RS2] C. M. Ringel, M. Schmidmeier, The Auslander-Reiten translation in submodule categories, Trans. Amer. Math. Soc. 360(2)(2008), 691-716.

[RS3] C. M. Ringel, M. Schmidmeier, Invariant subspaces of nilpotent operators I, J. rein angew. Math. 614(2008), 1-52.

[S] D. Simson, Representation types of the category of subprojective representations of a finite poset over $K[t]/(t^m)$ and a solution of a Birkhoff type problem, J. Algebra 311(2007), 1-30.

[SW] D. Simson, M. Wojewodzki, An algorithmic solution of a Birkhoff type problem, Fundamenta Informaticae 83(2008), 389-410.

[Z] P. Zhang, Monomorphism categories, cotilting theory, and Gorenstein-projective modules, preprint (2009). Avaible in arXiv: math. RT 1101.3872, 2011.

DEPARTMENT OF MATHEMATICS, SHANGHAI JIAO TONG UNIVERSITY, SHANGHAI 200240, P. R. CHINA

XIONGBAOLIN@GMAIL.COM, PZHANG@SJTU.EDU.CN, ZYH@SJTU.EDU.CN